

Anti-CXCL7 / PPBP Antibody

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS17299

Product Information

Application	WB, IHC-P
Primary Accession	P02775
Predicted	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	13894


Additional Information

Gene ID	5473
Alias Symbol Other Names	PPBP PPBP, Beta-thromboglobulin, B-TG1, Beta-TG, C-X-C motif chemokine 7, CTAP3, CXCL7, CTAP-III, CXC chemokine ligand 7, NAP-2, NAP-2-L1, PBP, LA-PF4, LDGF, SCYB7, Small-inducible cytokine B7, Small inducible cytokine B7, THBGB1, Thrombocidin 2, TC2, TGB ...
Target/Specificity	Human PPBP / CXCL7
Reconstitution & Storage	PBS, 0.1% sodium azide, 50% glycerol Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.
Precautions	Anti-CXCL7 / PPBP Antibody is for research use only and not for use in diagnostic or therapeutic procedures.


Protein Information

Name	PPBP
Synonyms	CTAP3, CXCL7, SCYB7, TGB1, THBGB1
Function	LA-PF4 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by human synovial cells. NAP-2 is a ligand for CXCR1 and CXCR2, and NAP-2, NAP-2(73), NAP-2(74), NAP-2(1-66), and most potent NAP-2(1-63) are chemoattractants and activators for neutrophils. TC-1 and TC-2 are antibacterial proteins, in vitro released from activated platelet alpha-granules. CTAP-III(1-81) is more potent than CTAP-III desensitize chemokine-induced neutrophil activation. Secreted.

Images



Human Placenta: Formalin-Fixed, Paraffin-Embedded (FFPE)



Human Spleen: Formalin-Fixed, Paraffin-Embedded (FFPE)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.