

Anti-MAD2L1 / MAD2 Antibody

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS17355

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q13257
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	23510

Additional Information

Gene ID	4085
Alias Symbol	MAD2L1
Other Names	MAD2L1, HSMAD2, MAD2, MAD2-like protein 1
Target/Specificity	Human MAD2L1 / MAD2
Reconstitution & Storage	PBS, pH 7.3, 0.02% sodium azide, 50% glycerol. Long term: -80°C; Short term: -20°C. Avoid freeze-thaw cycles.
Precautions	Anti-MAD2L1 / MAD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MAD2L1
Synonyms	MAD2
Function	Component of the mitotic checkpoint complex (MCC), which inhibits APC/C and delays chromosome segregation until all chromosomes are properly attached to the mitotic spindle (PubMed:10700282, PubMed:11804586, PubMed:15024386, PubMed:29162720, PubMed:27509861). The MCC, which consists of CDC20, MAD2L1/MAD2, BUB1B/BUBR1 and BUB3, is a key player of the spindle assembly checkpoint (PubMed:15020684, PubMed:27509861). The MCC-bound APC/C complex is inactive and this delays the metaphase/anaphase transition (PubMed:27509861). When the spindle assembly checkpoint is silenced, the MCC-APC/C complex recruits the E2 enzyme UBCH10 and triggers the auto-ubiquitination of CDC20 in the MCC, thereby reactivating APC/C for transition into anaphase (PubMed:27509861). In the closed conformation (C-MAD2) forms a heterotetrameric complex with

MAD1L1 at unattached kinetochores during prometaphase, the complex recruits open conformation molecules of MAD2L1 (O-MAD2) and then promotes the conversion of O-MAD2 to C-MAD2 (PubMed:[29162720](#)).

Cellular Location

Nucleus. Chromosome, centromere, kinetochore. Cytoplasm. Cytoplasm, cytoskeleton, spindle pole Note=Recruited by MAD1L1 to unattached kinetochores (Probable) Recruited to the nuclear pore complex by TPR during interphase Recruited to kinetochores in late prometaphase after BUB1, CENPF, BUB1B and CENPE. Kinetochore association requires the presence of NEK2 Kinetochore association is repressed by UBD. Sequestered to the cytoplasm upon interaction with isoform 3 of MAD1L1 (PubMed:19010891) {ECO:0000269|PubMed:19010891, ECO:0000305}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.