

Anti-IFNAR2 Antibody

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS17356

Product Information

Application	WB, IHC-P
Primary Accession	P48551
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	57759

Additional Information

Gene ID	3455
Alias Symbol	IFNAR2
Other Names	IFNAR2, IFN-alpha/beta receptor 2, IFN-alpha-REC, IFN-alpha binding protein, IFN-R, IFN-R-2, IFNARB, Type I interferon receptor 2, IFNABR
Target/Specificity	Human IFNAR2
Reconstitution & Storage	PBS, pH 7.3, 0.02% sodium azide, 50% glycerol. Long term: -80°C; Short term: -20°C. Avoid freeze-thaw cycles.
Precautions	Anti-IFNAR2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IFNAR2
Synonyms	IFNABR, IFNARB
Function	Together with IFNAR1, forms the heterodimeric receptor for type I interferons (including interferons alpha, beta, epsilon, omega and kappa) (PubMed: 10049744 , PubMed: 10556041 , PubMed: 21854986 , PubMed: 26424569 , PubMed: 28165510 , PubMed: 32972995 , PubMed: 7665574 , PubMed: 7759950 , PubMed: 8181059 , PubMed: 8798579 , PubMed: 8969169). Type I interferon binding activates the JAK-STAT signaling cascade, resulting in transcriptional activation or repression of interferon-regulated genes that encode the effectors of the interferon response (PubMed: 10049744 , PubMed: 17517919 , PubMed: 21854986 , PubMed: 26424569 , PubMed: 28165510 , PubMed: 32972995 , PubMed: 7665574 , PubMed: 7759950 , PubMed: 8181059 , PubMed: 8798579 , PubMed: 8969169). Mechanistically, type

I interferon-binding brings the IFNAR1 and IFNAR2 subunits into close proximity with one another, driving their associated Janus kinases (JAKs) (TYK2 bound to IFNAR1 and JAK1 bound to IFNAR2) to cross-phosphorylate one another (PubMed:[10556041](#), PubMed:[11682488](#), PubMed:[12105218](#), PubMed:[21854986](#), PubMed:[32972995](#)). The activated kinases phosphorylate specific tyrosine residues on the intracellular domains of IFNAR1 and IFNAR2, forming docking sites for the STAT transcription factors (STAT1, STAT2 and STAT3) (PubMed:[11682488](#), PubMed:[12105218](#), PubMed:[21854986](#), PubMed:[32972995](#)). STAT proteins are then phosphorylated by the JAKs, promoting their translocation into the nucleus to regulate expression of interferon-regulated genes (PubMed:[12105218](#), PubMed:[28165510](#), PubMed:[9121453](#)).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Isoform 3]: Secreted

Tissue Location

Isoform 3 is detected in the urine (at protein level) (PubMed:7759950, PubMed:8181059). Expressed in blood cells Expressed in lymphoblastoid and fibrosarcoma cell lines

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.