

Anti-AIMP1 / EMAP II Antibody

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS17362

Product Information

Application	WB, IHC-P, IF
Primary Accession	Q12904
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	34353

Additional Information

Gene ID	9255
Alias Symbol	AIMP1
Other Names	AIMP1, EMAP II, EMAPII, EMAP-2, HLD3, p43, EMAP2, SCYE1
Target/Specificity	Human AIMP1 / EMAP II / p43
Reconstitution & Storage	PBS, pH 7.3, 0.02% sodium azide, 50% glycerol. Long term: -80°C; Short term: -20°C. Avoid freeze-thaw cycles.
Precautions	Anti-AIMP1 / EMAP II Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	AIMP1
Synonyms	EMAP2, SCYE1
Function	Non-catalytic component of the multisynthase complex. Stimulates the catalytic activity of cytoplasmic arginyl-tRNA synthase (PubMed: 10358004). Binds tRNA. Possesses inflammatory cytokine activity (PubMed: 11306575). Negatively regulates TGF-beta signaling through stabilization of SMURF2 by binding to SMURF2 and inhibiting its SMAD7- mediated degradation (By similarity). Involved in glucose homeostasis through induction of glucagon secretion at low glucose levels (By similarity). Promotes dermal fibroblast proliferation and wound repair (PubMed: 16472771). Regulates KDELR1-mediated retention of HSP90B1/gp96 in the endoplasmic reticulum (By similarity). Plays a role in angiogenesis by inducing endothelial cell migration at low concentrations and endothelial cell apoptosis at high concentrations (PubMed: 12237313). Induces maturation of dendritic cells and

monocyte cell adhesion (PubMed:[11818442](#)). Modulates endothelial cell responses by degrading HIF-1A through interaction with PSMA7 (PubMed:[19362550](#)).

Cellular Location

Nucleus. Cytoplasm, cytosol. Secreted. Endoplasmic reticulum {ECO:0000250|UniProtKB:P31230}. Golgi apparatus {ECO:0000250|UniProtKB:P31230}. Note=Enriched in secretory vesicles of pancreatic alpha cells and secreted from the pancreas in response to low glucose levels (By similarity). Secreted in response to hypoxia (PubMed:10850427). Also secreted in response to both apoptotic and necrotic cell death. {ECO:0000250|UniProtKB:P31230, ECO:0000269|PubMed:10850427}

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.