10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-SFN / Stratifin / 14-3-3 Sigma Antibody

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17392

Product Information

Application	WB, IHC-P
Primary Accession	P31947
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27774
Concentration (mg/ml)	0.5 mg/ml

Additional Information

Gene ID	2810
Alias Symbol	SFN
Other Names	SFN, 14-3-3 sigma, 14-3-3 protein sigma, HME1, Stratifin, YWHAS
Target/Specificity	Human SFN / Stratifin / 14-3-3 Sigma
Reconstitution & Storage	PBS, 0.05% sodium azide. Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.
Precautions	Anti-SFN / Stratifin / 14-3-3 Sigma Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SFN (HGNC:10773)
Function	Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed:15731107, PubMed:22634725, PubMed:28202711, PubMed:37797010). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed:15731107, PubMed:22634725, PubMed:28202711, PubMed:37797010). Binding generally results in the modulation of the activity of the binding partner (PubMed:15731107, PubMed:22634725, PubMed:28202711, PubMed:37797010). Promotes cytosolic retention of GBP1 GTPase by binding to phosphorylated GBP1, thereby inhibiting the innate immune response (PubMed:37797010). Also acts as a TP53/p53-regulated inhibitor of G2/M progression (PubMed:9659898). When bound to KRT17, regulates protein synthesis and epithelial cell growth by stimulating Akt/mTOR pathway (By similarity). Acts to maintain desmosome cell junction adhesion in epithelial cells via interacting with and

	sequestering PKP3 to the cytoplasm, thereby restricting its translocation to existing desmosome structures and therefore maintaining desmosome protein homeostasis (PubMed:24124604). Also acts to facilitate PKP3 exchange at desmosome plaques, thereby maintaining keratinocyte intercellular adhesion (PubMed:29678907). May also regulate MDM2 autoubiquitination and degradation and thereby activate p53/TP53 (PubMed:18382127). Upstream regulator of YAP1 localization in keratinocytes: binds and sequester YAP1 in the cytoplasm, limiting YAP1-dependent transcription and driving Hippo signaling and cell differentiation (By similarity).
Cellular Location	Cytoplasm. Nucleus. Secreted Note=May be secreted by a non-classical secretory pathway
Tissue Location	Present mainly in tissues enriched in stratified squamous keratinizing epithelium.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.