10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-PRIM1 Antibody (C-Terminus)

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17410

Product Information

Application	WB, IHC-P
Primary Accession	P49642
Predicted	Human, Mouse, Rabbit, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49902
Concentration (mg/ml)	1 mg/ml

Additional Information

Gene ID	5557
Alias Symbol	PRIM1
Other Names	PRIM1, DNA primase subunit 48, DNA primase 1, Dna primase 49 kd subunit, DNA primase 49 kDa subunit, p49, Primase polypeptide 1, 49kDa, DNA primase small subunit, Primase p49 subunit
Target/Specificity	Recognizes endogenous levels of PRIM1 protein.
Reconstitution & Storage	PBS, pH 7.3, 0.01% sodium azide, 30% glycerol. Store at -20°C. Aliquot to avoid freeze/thaw cycles.
Precautions	Anti-PRIM1 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PRIM1
Function	Catalytic subunit of the DNA primase complex and component of the DNA polymerase alpha complex (also known as the alpha DNA polymerase-primase complex - primosome/replisome) which play an essential role in the initiation of DNA synthesis (PubMed:17893144, PubMed:24043831, PubMed:25550159, PubMed:26975377, PubMed:31479243, PubMed:33060134, PubMed:9268648, PubMed:9705292). During the S phase of the cell cycle, the DNA polymerase alpha complex (composed of a catalytic subunit POLA1, an accessory subunit POLA2 and two primase subunits, the catalytic subunit PRIM1 and the regulatory subunit PRIM2) is recruited to DNA at the replicative forks via direct interactions with MCM10 and WDHD1 (By similarity). The primase subunit of the polymerase alpha complex initiates DNA synthesis by oligomerizing short RNA primers on both leading and

lagging strands (PubMed:17893144). These primers are initially extended by the polymerase alpha catalytic subunit and subsequently transferred to polymerase delta and polymerase epsilon for processive synthesis on the lagging and leading strand, respectively (By similarity). In the primase complex, both subunits are necessary for the initial di-nucleotide formation, but the extension of the primer depends only on the catalytic subunit (PubMed:17893144). Synthesizes 9-mer RNA primers (also known as the 'unit length' RNA primers). Incorporates only ribonucleotides in the presence of ribo- and deoxy-nucleotide triphosphates (rNTPs, dNTPs) (PubMed:26975377). Requires template thymine or cytidine to start the RNA primer synthesis, with an adenine or guanine at its 5'-end (PubMed:25550159, PubMed:26975377). Binds single stranded DNA (By similarity).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.