10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-HELO1 / ELOVL5 Antibody (N-Terminus)

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17680

Product Information

Application	IHC-P
Primary Accession	Q9NYP7
Predicted	Human, Monkey, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35293
Concentration (mg/ml)	1 mg/ml

Additional Information

Gene ID	60481
Alias Symbol	ELOVL5
Other Names	ELOVL5, ELOVL FA elongase 5, ELOVL fatty acid elongase 5, Fatty acid elongase 1, RP3-483K16.1, DJ483K16.1, HELO1
Target/Specificity	Human ELOVL5. BLAST analysis of the peptide immunogen showed no homology with other human proteins.
Reconstitution & Storage	Immunoaffinity purified
Precautions	Anti-HELO1 / ELOVL5 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ELOVL5 {ECO:0000255|HAMAP-Rule:MF_03205}
Synonyms	ELOVL2
Function	Catalyzes the first and rate-limiting reaction of the four reactions that constitute the long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process allows the addition of 2 carbons to the chain of long- and very long-chain fatty acids (VLCFAs) per cycle. Condensing enzyme that acts specifically toward polyunsaturated acyl-CoA with the higher activity toward C18:3(n-6) acyl-CoA. May participate in the production of monounsaturated and of polyunsaturated VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators (By similarity) (PubMed:10970790, PubMed:20937905). In conditions where the essential linoleic and alpha linoleic fatty acids are lacking it is also involved in the synthesis of Mead acid

	from oleic acid (By similarity).
Cellular Location	Endoplasmic reticulum membrane {ECO:0000255|HAMAP-Rule:MF_03205, ECO:0000269|PubMed:20937905}; Multi- pass membrane protein {ECO:0000255|HAMAP-Rule:MF_03205}. Cell projection, dendrite {ECO:0000255|HAMAP-Rule:MF_03205, ECO:0000269|PubMed:25065913}. Note=In Purkinje cells, the protein localizes to the soma and proximal portion of the dendritic tree {ECO:0000255|HAMAP-Rule:MF_03205, ECO:0000269|PubMed:25065913}
Tissue Location	Ubiquitous. Highly expressed in the adrenal gland and testis. Weakly expressed in prostate, lung and brain. Expressed in the cerebellum.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.