

Anti-IGF1 Antibody

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS17732

Product Information

Application	WB, IHC-P, E, Neut
Primary Accession	P05019
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	21841

Additional Information

Gene ID	3479
Alias Symbol	IGF1
Other Names	IGF1, IGF-IA, IGF-IB, IGFI, IBP1, IGF-I, Insulin-like growth factor 1, Insulin-like growth factor I, Insulin-like growth factor IA, Insulin-like growth factor IB, Mechano growth factor, Somatomedin-C, IGF1A, MGF
Target/Specificity	Human IGF-I
Reconstitution & Storage	Immunoaffinity purified
Precautions	Anti-IGF1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	IGF1 (HGNC:5464)
Function	The insulin-like growth factors, isolated from plasma, are structurally and functionally related to insulin but have a much higher growth-promoting activity. May be a physiological regulator of [1-14C]- 2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to enhancing glucose uptake. May play a role in synapse maturation (PubMed: 21076856 , PubMed: 24132240). Ca(2+)-dependent exocytosis of IGF1 is required for sensory perception of smell in the olfactory bulb (By similarity). Acts as a ligand for IGF1R. Binds to the alpha subunit of IGF1R, leading to the activation of the intrinsic tyrosine kinase activity which autophosphorylates tyrosine residues in the beta subunit thus initiating a cascade of down-stream signaling events leading to activation of the

PI3K-AKT/PKB and the Ras-MAPK pathways. Binds to integrins ITGA5:ITGB3 and ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and IGFR1 are essential for IGF1 signaling. Induces the phosphorylation and activation of IGFR1, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed:[19578119](#), PubMed:[22351760](#), PubMed:[23243309](#), PubMed:[23696648](#)). As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via promotion of STUB1/CHIP-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity).

Cellular Location

Secreted {ECO:0000250|UniProtKB:P05017}.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.