10320 Camino Santa Fe, Suite G San Diego, CA 92121 Tel: 858.875.1900 Fax: 858.875.1999



## Anti-CD19 Antibody (C-Terminus)

Goat Anti Human Polyclonal Antibody Catalog # ALS18239

## **Product Information**

Application WB, IHC-P Primary Accession P15391

**Predicted** Human, Mouse, Rat, Monkey, Dog

Host Goat
Clonality Polyclonal
Isotype IgG
Calculated MW 61128
Concentration (mg/ml) 3 mg/ml

## **Additional Information**

Gene ID 930

Alias Symbol CD19

Other Names CD19, B-lymphocyte antigen CD19, CD19 molecule, Differentiation antigen

CD19, T-cell surface antigen Leu-12, B4, CD19 antigen, CVID3

**Target/Specificity** Using spleen and Jurkat cell lysates detects a 95 kDa band by Western blot.

Reconstitution & Storage Immunoaffinity purified

**Precautions** Anti-CD19 Antibody (C-Terminus) is for research use only and not for use in

diagnostic or therapeutic procedures.

## **Protein Information**

Name CD19

**Function** Functions as a coreceptor for the B-cell antigen receptor complex (BCR) on

B-lymphocytes (PubMed: <u>29523808</u>). Decreases the threshold for activation of downstream signaling pathways and for triggering B-cell responses to antigens (PubMed: <u>1373518</u>, PubMed: <u>16672701</u>, PubMed: <u>2463100</u>). Activates signaling pathways that lead to the activation of phosphatidylinositol 3-kinase and the mobilization of intracellular Ca(2+) stores (PubMed: <u>12387743</u>,

PubMed:<u>16672701</u>, PubMed:<u>9317126</u>, PubMed:<u>9382888</u>). Is not required for early steps during B cell differentiation in the blood marrow

(PubMed:<u>9317126</u>). Required for normal differentiation of B-1 cells (By similarity). Required for normal B cell differentiation and proliferation in response to antigen challenges (PubMed:<u>1373518</u>, PubMed:<u>2463100</u>).

Required for normal levels of serum immunoglobulins, and for production of high-affinity antibodies in response to antigen challenge (PubMed: 12387743,

PubMed: 16672701, PubMed: 9317126).

**Cellular Location** Cell membrane; Single-pass type I membrane protein. Membrane raft

{ECO:0000250|UniProtKB:P25918}; Single-pass type I membrane protein

{ECO:0000250 | UniProtKB:P25918}

**Tissue Location** Detected on marginal zone and germinal center B cells in lymph nodes

(PubMed:2463100). Detected on blood B cells (at protein level)

(PubMed:16672701, PubMed:2463100)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.