

# Anti-GRN / Granulin Antibody

Rabbit Anti Human Polyclonal Antibody  
Catalog # ALS18342

## Product Information

---

<b>Application</b>	WB, IHC-P
<b>Primary Accession</b>	<a href="#">P28799</a>
<b>Predicted</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Calculated MW</b>	63544

## Additional Information

---

<b>Gene ID</b>	2896
<b>Alias Symbol</b>	GRN
<b>Other Names</b>	GRN, Acrogranin, CLN11, Epithelin precursor, GEP, gp88, Granulin, Granulin-epithelin, Granulins, PC cell-derived growth factor, PCDGF, PEPI, PGRN, Proepithelin, Progranulin
<b>Target/Specificity</b>	Human GRN / Granulin.
<b>Reconstitution &amp; Storage</b>	Affinity purified
<b>Precautions</b>	Anti-GRN / Granulin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	GRN ( <a href="#">HGNC:4601</a> )
<b>Function</b>	Secreted protein that acts as a key regulator of lysosomal function and as a growth factor involved in inflammation, wound healing and cell proliferation (PubMed: <a href="#">12526812</a> , PubMed: <a href="#">18378771</a> , PubMed: <a href="#">28073925</a> , PubMed: <a href="#">28453791</a> , PubMed: <a href="#">28541286</a> ). Regulates protein trafficking to lysosomes, and also the activity of lysosomal enzymes (PubMed: <a href="#">28453791</a> , PubMed: <a href="#">28541286</a> ). Also facilitates the acidification of lysosomes, causing degradation of mature CTSD by CTSB (PubMed: <a href="#">28073925</a> ). In addition, functions as a wound-related growth factor that acts directly on dermal fibroblasts and endothelial cells to promote division, migration and the formation of capillary-like tubule structures (By similarity). Also promotes epithelial cell proliferation by blocking TNF-mediated neutrophil activation preventing release of oxidants and proteases (PubMed: <a href="#">12526812</a> ). Moreover, modulates inflammation in neurons by preserving neurons survival, axonal

outgrowth and neuronal integrity (PubMed:[18378771](#)).

**Cellular Location**

Secreted. Lysosome Note=Endocytosed by SORT1 and delivered to lysosomes (PubMed:21092856, PubMed:28073925). Targeted to lysosome by PSAP via M6PR and LRP1, in both biosynthetic and endocytic pathways (PubMed:26370502, PubMed:28073925). Co-localized with GBA1 in the intracellular trafficking compartments until to lysosome (By similarity) {ECO:0000250|UniProtKB:P28798, ECO:0000269|PubMed:21092856, ECO:0000269|PubMed:26370502, ECO:0000269|PubMed:28073925}

**Tissue Location**

In myelogenous leukemic cell lines of promonocytic, promyelocytic, and proerythroid lineage, in fibroblasts, and very strongly in epithelial cell lines. Present in inflammatory cells and bone marrow. Highest levels in kidney

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.