10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-ABCC10 Antibody (aa835-884)

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18362

Product Information

Application	WB, IHC-P, E
Primary Accession	Q5T3U5
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	161629

Additional Information

Gene ID	89845
Alias Symbol	ABCC10
Other Names	ABCC10, EST182763, Multidrug resistance protein 7, MRP7, SIMRP7
Target/Specificity	MRP7 Antibody detects endogenous levels of total MRP7 protein.
Reconstitution & Storage	Immunoaffinity purified
Precautions	Anti-ABCC10 Antibody (aa835-884) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ABCC10 {ECO:0000303|PubMed:12527806}
Function	ATP-dependent transporter of the ATP-binding cassette (ABC) family that actively extrudes physiological compounds, and xenobiotics from cells. Lipophilic anion transporter that mediates ATP-dependent transport of glucuronide conjugates such as estradiol-17-beta-o- glucuronide and GSH conjugates such as leukotriene C4 (LTC4) (PubMed:12527806, PubMed:15256465). May regulate the transport of organic compounds in testes across the blood-testis-barrier (Probable). Mediates intercellular propagation of antiviral immune signaling in early stages of infection (By similarity). In RNA virus-infected cells, oligoadenylate synthase senses viral dsRNA and generates 2',5'- oligoadenylates (2-5A) which act as second messengers to activate RNASEL and type I interferon signaling to inhibit viral replication (By similarity). This innate signaling pathway is locally extended and amplified by ABCC10, which exports 2-5A from virus-infected cells to cross-activates RNASEL in uninfected neighboring cells and confers protection against viral infection (By similarity). Mediates multidrug resistance (MDR) in

	cancer cells by preventing the intracellular accumulation of certain antitumor drugs, such as, docetaxel and paclitaxel (PubMed:15256465, PubMed:23087055). Does not transport glycocholic acid, taurocholic acid, MTX, folic acid, cAMP, or cGMP (PubMed:12527806).
Cellular Location	Cell membrane; Multi-pass membrane protein {ECO:0000255|PROSITE-ProRule:PRU00441, ECO:0000269|PubMed:12566991, ECO:0000269|PubMed:15256465}. Basolateral cell membrane; Multi-pass membrane protein. Basal cell membrane; Multi-pass membrane protein. Note=Localized to the basal membrane of Sertoli cells.
Tissue Location	In testis, localized to peritubular myoid cells, Leydig cells, along the basal membrane of Sertoli cells, moderately in the adluminal compartment of the seminiferous tubules, and in vascular endothelial cells. [Isoform 2]: Widely expressed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.