

Anti-CYP2D6 Antibody (C-Terminus)

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS18410

Product Information

Application	WB, IHC-P
Primary Accession	P10635
Predicted	Human, Mouse, Rat, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55769
Concentration (mg/ml)	1 mg/ml

Additional Information

Gene ID	1565
Alias Symbol Other Names	CYP2D6 CYP2D6, CPD6, CYP11D6, Cytochrome P450 2D6, CYP2D, CYP2D8P2, Cytochrome p450 iid6, CYP2D7P2, CYP2DL1, Cytochrome P450-DB1, Debrisoquine 4-hydroxylase, p450C2D, p450-DB1, p450DB1, CYP2D7AP, CYP2D7BP
Target/Specificity	Recognizes endogenous levels of Cytochrome P450 2D6 protein.
Reconstitution & Storage	Immunoaffinity purified
Precautions	Anti-CYP2D6 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CYP2D6 {ECO:0000303 PubMed:21289075, ECO:0000312 HGNC:HGNC:2625}
Function	A cytochrome P450 monooxygenase involved in the metabolism of fatty acids, steroids and retinoids (PubMed: 18698000 , PubMed: 19965576 , PubMed: 20972997 , PubMed: 21289075 , PubMed: 21576599). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed: 18698000 , PubMed: 19965576 , PubMed: 20972997 , PubMed: 21289075 , PubMed: 21576599). Catalyzes the epoxidation of double bonds of polyunsaturated fatty acids (PUFA) (PubMed: 19965576 , PubMed: 20972997). Metabolizes endocannabinoid arachidonylethanolamide (anandamide) to 20-hydroxyeicosatetraenoic acid ethanolamide (20-HETE-EA)

and 8,9-, 11,12-, and 14,15-epoxyeicosatrienoic acid ethanolamides (EpETrE-EAs), potentially modulating endocannabinoid system signaling (PubMed:[18698000](#), PubMed:[21289075](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Metabolizes cholesterol toward 25-hydroxycholesterol, a physiological regulator of cellular cholesterol homeostasis (PubMed:[21576599](#)). Catalyzes the oxidative transformations of all-trans retinol to all-trans retinal, a precursor for the active form all-trans-retinoic acid (PubMed:[10681376](#)). Also involved in the oxidative metabolism of drugs such as antiarrhythmics, adrenoceptor antagonists, and tricyclic antidepressants.

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.