

Anti-NMNAT2 Antibody (clone Nady-1)

Mouse Anti Human Monoclonal Antibody

Catalog # ALS18499

Product Information

Application	WB, IHC-P, E
Primary Accession	Q9BZQ4
Predicted	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,k
Clone Names	Nady-1
Calculated MW	34439
Concentration (mg/ml)	1 mg/ml

Additional Information

Gene ID	23057
Alias Symbol	NMNAT2
Other Names	NMNAT2, C1orf15, KIAA0479, NaMN adenylyltransferase 2, NMN adenylyltransferase 2, PNAT2
Target/Specificity	Recognizes human NMNAT2.
Reconstitution & Storage	Protein G purified
Precautions	Anti-NMNAT2 Antibody (clone Nady-1) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	NMNAT2 (HGNC:16789)
Function	Nicotinamide/nicotinate-nucleotide adenylyltransferase that acts as an axon maintenance factor (By similarity). Axon survival factor required for the maintenance of healthy axons: acts by delaying Wallerian axon degeneration, an evolutionarily conserved process that drives the loss of damaged axons (By similarity). Catalyzes the formation of NAD(+) from nicotinamide mononucleotide (NMN) and ATP (PubMed: 16118205 , PubMed: 17402747). Can also use the deamidated form; nicotinic acid mononucleotide (NaMN) as substrate but with a lower efficiency (PubMed: 16118205 , PubMed: 17402747). Cannot use triazofurin monophosphate (TrMP) as substrate (PubMed: 16118205 , PubMed: 17402747). Also catalyzes the reverse reaction, i.e. the pyrophosphorolytic cleavage of NAD(+) (PubMed: 16118205 , PubMed: 17402747). For the pyrophosphorolytic activity prefers NAD(+), NADH

and NaAD as substrates and degrades nicotinic acid adenine dinucleotide phosphate (NAD) less effectively (PubMed:[16118205](#), PubMed:[17402747](#)). Fails to cleave phosphorylated dinucleotides NADP(+), NADPH and NaADP(+) (PubMed:[16118205](#), PubMed:[17402747](#)). Also acts as an activator of ADP-ribosylation by supporting the catalytic activity of PARP16 and promoting mono-ADP-ribosylation of ribosomes by PARP16 (PubMed:[34314702](#)). May be involved in the maintenance of axonal integrity (By similarity).

Cellular Location

Golgi apparatus membrane; Lipid-anchor {ECO:0000250|UniProtKB:Q8BNJ3}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q8BNJ3}; Lipid-anchor {ECO:0000250|UniProtKB:Q8BNJ3}. Cytoplasm. Cell projection, axon {ECO:0000250|UniProtKB:Q8BNJ3}. Note=Delivered to axons with Golgi-derived cytoplasmic vesicles. {ECO:0000250|UniProtKB:Q8BNJ3}

Tissue Location

Highly expressed in brain, in particular in cerebrum, cerebellum, occipital lobe, frontal lobe, temporal lobe and putamen. Also found in the heart, skeletal muscle, pancreas and islets of Langerhans.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.