

Anti-ATG3 Antibody (aa1-314)

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS18512

Product Information

Application	WB, IHC-P, E
Primary Accession	Q9NT62
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	35864

Additional Information

Gene ID	64422
Alias Symbol	ATG3
Other Names	ATG3, APG3-LIKE, APG3, APG3L, Autophagy related 3, HAPg3, PC3-96, Protein PC3-96, Autophagy-related protein 3
Target/Specificity	Human ATG3
Reconstitution & Storage	Caprylic acid and ammonium sulfate precipitation
Precautions	Anti-ATG3 Antibody (aa1-314) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ATG3 (HGNC:20962)
Synonyms	APG3, APG3L
Function	E2 conjugating enzyme that catalyzes the covalent conjugation of the C-terminal Gly of ATG8-like proteins (GABARAP, GABARAPL1, GABARAPL2 or MAP1LC3A) to the amino group of phosphatidylethanolamine (PE)-containing lipids in the membrane resulting in membrane-bound ATG8-like proteins which is one of the key steps in the development of autophagic isolation membranes during autophagosome formation (PubMed: 24191030 , PubMed: 33446636 , PubMed: 37252361). Cycles back and forth between binding to ATG7 for loading with the ATG8-like proteins and binding to E3 enzyme, composed of ATG12, ATG5 and ATG16L1 to promote ATG8-like proteins lipidation (PubMed: 11825910 , PubMed: 12207896 , PubMed: 12890687 , PubMed: 16704426 , PubMed: 24186333). Also plays a role as a membrane curvature sensor that facilitates LC3/GABARAP lipidation by

sensing local membrane stress associated with lipid-packing defects as occurs with high molar proportions of conical lipids or strident membrane curvature (By similarity). Interacts with negatively-charged membranes promoting membrane tethering and enhancing LC3/GABARAP lipidation (PubMed:[29142222](#)). Also acts as an autocatalytic E2-like enzyme by catalyzing the conjugation of ATG12 to itself in an ATG7-dependent manner, this complex thus formed, plays a role in mitochondrial homeostasis but not in autophagy (By similarity). ATG12- ATG3 conjugation promotes late endosome to lysosome trafficking and basal autophagosome maturation via its interaction with PDCD6IP (By similarity). ATG12-ATG3 conjugate is also formed upon vaccinia virus infection, leading to the disruption the cellular autophagy which is not necessary for vaccinia survival and proliferation (By similarity). Promotes primary ciliogenesis by removing OFD1 from centriolar satellites via the autophagic pathway (By similarity).

Cellular Location

Cytoplasm.

Tissue Location

Widely expressed, with a highest expression in heart, skeletal muscle, kidney, liver and placenta

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.