

Anti-DDX39B / UAP56 Antibody (aa2-251)

Rabbit Anti Human Polyclonal Antibody

Catalog # ALS18515

Product Information

Application	WB, IHC-P, E
Primary Accession	Q13838
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	48991

Additional Information

Gene ID	7919
Alias Symbol	DDX39B
Other Names	DDX39B, BAT1, DEAD box protein UAP56, HLA-B associated transcript 1, UAP56, ATP-dependent RNA helicase p47, D6S81E, Spliceosome RNA helicase BAT1
Target/Specificity	Human DDX39B / UAP56 / BAT1
Reconstitution & Storage	Caprylic acid and ammonium sulfate precipitation
Precautions	Anti-DDX39B / UAP56 Antibody (aa2-251) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	DDX39B (HGNC:13917)
Synonyms	BAT1, UAP56
Function	Involved in nuclear export of spliced and unspliced mRNA (PubMed: 15833825 , PubMed: 15998806 , PubMed: 17190602). Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA (PubMed: 15833825 , PubMed: 15998806 , PubMed: 17190602). The TREX complex is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap- dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NXF1 pathway (PubMed: 15833825 , PubMed: 15998806 , PubMed: 17190602). The THOC1-THOC2- THOC3 core

complex alone is sufficient to promote ATPase activity of DDX39B; in the complex THOC2 is the only component that directly interacts with DDX39B (PubMed:[33191911](#)). Associates with SARNP/CIP29, which facilitates RNA binding of DDX39B and likely plays a role in mRNA export (PubMed:[37578863](#)). May undergo several rounds of ATP hydrolysis during assembly of TREX to drive subsequent loading of components such as ALYREF/THOC4 and CHTOP onto mRNA. Also associates with pre-mRNA independent of ALYREF/THOC4. Involved in the nuclear export of intronless mRNA; the ATP-bound form is proposed to recruit export adapter ALYREF/THOC4 to intronless mRNA; its ATPase activity is cooperatively stimulated by RNA and ALYREF/THOC4 and ATP hydrolysis is thought to trigger the dissociation from RNA to allow the association of ALYREF/THOC4 and the NXF1-NXT1 heterodimer. Involved in transcription elongation and genome stability.

Cellular Location

Nucleus. Nucleus speckle. Cytoplasm. Note=Can translocate to the cytoplasm in the presence of MX1. TREX complex assembly seems to occur in regions surrounding nuclear speckles known as perispeckles

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.