

## Anti-CCND2 / Cyclin D2 Antibody

Rabbit Anti Human Polyclonal Antibody Catalog # ALS18584

## **Product Information**

Application	WB, IHC-P, IF
Primary Accession	<u>P30279</u>
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	33067

## **Additional Information**

Gene ID	894
Alias Symbol Other Names	CCND2 CCND2, KIAK0002, G1/S-specific cyclin D2, G1/S-specific cyclin-D2, Cyclin D2
Target/Specificity	Human CCND2 / Cyclin D2
Reconstitution & Storage	Affinity purified
Precautions	Anti-CCND2 / Cyclin D2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **Protein Information**

Name	CCND2 {ECO:0000303 PubMed:1386336, ECO:0000312 HGNC:HGNC:1583}
Function	Regulatory component of the cyclin D2-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed: <u>18827403</u> , PubMed: <u>8114739</u> ). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed: <u>18827403</u> , PubMed: <u>8114739</u> ). Hypophosphorylates RB1 in early G(1) phase (PubMed: <u>18827403</u> , PubMed: <u>8114739</u> ). Cyclin D-CDK4 complexes are major integrators of various mitogenenic and antimitogenic signals (PubMed: <u>18827403</u> , PubMed: <u>8114739</u> ).
Cellular Location	Nucleus. Cytoplasm. Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated into the nucleus through interaction with KIP/CIP family members

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.