

Anti-RBX1 / ROC1 Antibody

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18585

Product Information

Application	WB, IHC-P, IF
Primary Accession	P62877
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	12274

Additional Information

Gene ID	9978
Alias Symbol Other Names	RBX1 RBX1, Regulator of cullins 1, RING box protein 1, ROC1, TESTICAN, Ring-box 1, RNF75, BA554C12.1, Protein ZYP, RING finger protein 75, RING-box protein 1, ZYP protein
Target/Specificity	Human ROC1 / RBX1
Reconstitution & Storage	Affinity purified
Precautions	Anti-RBX1 / ROC1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RBX1 (HGNC:9928)
Function	E3 ubiquitin ligase component of multiple cullin-RING-based E3 ubiquitin-protein ligase (CRLs) complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins, including proteins involved in cell cycle progression, signal transduction, transcription and transcription-coupled nucleotide excision repair (PubMed: 10230407 , PubMed: 10579999 , PubMed: 11961546 , PubMed: 15983046 , PubMed: 16678110 , PubMed: 19112177 , PubMed: 19679664 , PubMed: 22748924 , PubMed: 23455478 , PubMed: 27565346 , PubMed: 29769719 , PubMed: 32355176 , PubMed: 33417871 , PubMed: 37844242 , PubMed: 38326650 , PubMed: 39504960 , PubMed: 39667934 , PubMed: 38316879). CRLs complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins, ARIH1 mediating addition of the first ubiquitin on CRLs targets (PubMed: 27565346).

The functional specificity of the E3 ubiquitin-protein ligase complexes depends on the variable substrate recognition components. As a component of the CSA complex mediates ubiquitination of Pol II subunit POLR2A at 'Lys-1268', a critical TC- NER checkpoint (PubMed:[32355176](#), PubMed:[34526721](#)). Core component of the Cul7-RING(FBXW8) ubiquitin ligase complex, which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:[35982156](#)). Core component of a Cul9-RING ubiquitin ligase complex composed of CUL9 and RBX1, which mediates mono- ubiquitination of p53/TP53 (PubMed:[38605244](#)). Recruits the E2 ubiquitin-conjugating enzyme CDC34 to the complex and brings it into close proximity to the substrate. Probably also stimulates CDC34 autoubiquitination. May be required for histone H3 and histone H4 ubiquitination in response to ultraviolet and for subsequent DNA repair. Promotes the neddylation of CUL1, CUL2, CUL4 and CUL4 via its interaction with UBE2M. Involved in the ubiquitination of KEAP1, ENC1 and KLHL41. In concert with ATF2 and CUL3, promotes degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. As part of a multisubunit complex composed of elongin BC complex (ELOB and ELOC), elongin A/ELOA, RBX1 and CUL5; polyubiquitinates monoubiquitinated POLR2A (PubMed:[19920177](#)).

Cellular Location Cytoplasm. Nucleus

Tissue Location Widely expressed.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.