

Anti-SYVN1 / HRD1 Antibody

Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18596

Product Information

Application	WB, IHC-P
Primary Accession	Q86TM6
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	67685

Additional Information

Gene ID	84447
Alias Symbol	SYVN1
Other Names	SYVN1, KIAA1810, Synoviolin, Synovial apoptosis inhibitor 1, DER3, HRD1
Target/Specificity	Human SYVN1 / HRD1
Reconstitution & Storage	Affinity purified
Precautions	Anti-SYVN1 / HRD1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	SYVN1 {ECO:0000303 PubMed:15489334}
Function	E3 ubiquitin-protein ligase which accepts ubiquitin specifically from endoplasmic reticulum-associated UBC7 E2 ligase and transfers it to substrates, promoting their degradation (PubMed: 12459480 , PubMed: 12646171 , PubMed: 12975321 , PubMed: 14593114 , PubMed: 16289116 , PubMed: 16847254 , PubMed: 17059562 , PubMed: 17141218 , PubMed: 17170702 , PubMed: 22607976 , PubMed: 27827840 , PubMed: 26471130 , PubMed: 28827405). Component of the endoplasmic reticulum quality control (ERQC) system also called ER-associated degradation (ERAD) involved in ubiquitin-dependent degradation of misfolded endoplasmic reticulum proteins (PubMed: 12459480 , PubMed: 12646171 , PubMed: 12975321 , PubMed: 14593114 , PubMed: 16289116 , PubMed: 16847254 , PubMed: 17059562 , PubMed: 17141218 , PubMed: 17170702 , PubMed: 22607976 , PubMed: 26471130 , PubMed: 28842558). Also promotes the degradation of normal but naturally short-lived proteins such as SGK.

Protects cells from ER stress-induced apoptosis. Protects neurons from apoptosis induced by polyglutamine- expanded huntingtin (HTT) or unfolded GPR37 by promoting their degradation (PubMed:[17141218](#)). Sequesters p53/TP53 in the cytoplasm and promotes its degradation, thereby negatively regulating its biological function in transcription, cell cycle regulation and apoptosis (PubMed:[17170702](#)). Mediates the ubiquitination and subsequent degradation of cytoplasmic NFE2L1 (By similarity). During the early stage of B cell development, required for degradation of the pre-B cell receptor (pre-BCR) complex, hence supporting further differentiation into mature B cells (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein

Tissue Location

Ubiquitously expressed, with highest levels in liver and kidney (at protein level). Up-regulated in synovial tissues from patients with rheumatoid arthritis (at protein level)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.