

# Anti-EED Antibody

Rabbit Anti Human Polyclonal Antibody Catalog # ALS18600

#### **Product Information**

**Application** WB, IHC-P **Primary Accession** 075530

**Predicted** Human, Mouse, Rat

HostRabbitClonalityPolyclonalIsotypeIgGCalculated MW50198

# **Additional Information**

**Gene ID** 8726

Alias Symbol EED

Other Names EED, Embryonic ectoderm development, HEED, WAIT1, WAIT-1, Polycomb

protein EED

Target/Specificity Human EED

**Reconstitution & Storage** Affinity purified

**Precautions** Anti-EED Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

## **Protein Information**

Name EED ( HGNC:3188)

**Function** Polycomb group (PcG) protein. Component of the PRC2/EED-EZH2 complex,

which methylates 'Lys-9' and 'Lys-27' of histone H3, leading to transcriptional repression of the affected target gene. Also recognizes 'Lys-26' trimethylated histone H1 with the effect of inhibiting PRC2 complex methyltransferase activity on nucleosomal histone H3 'Lys-27', whereas H3 'Lys-27' recognition has the opposite effect, enabling the propagation of this repressive mark. The PRC2/EED- EZH2 complex may also serve as a recruiting platform for DNA methyltransferases, thereby linking two epigenetic repression systems. Genes repressed by the PRC2/EED-EZH2 complex include HOXC8, HOXA9, MYT1 and

CDKN2A.

**Cellular Location** Nucleus. Chromosome. Note=Transiently colocalizes with XIST at inactive X

chromosomes

## **Tissue Location**

Expressed in brain, colon, heart, kidney, liver, lung, muscle, ovary, peripheral blood leukocytes, pancreas, placenta, prostate, spleen, small intestine, testis, thymus and uterus. Appears to be overexpressed in breast and colon cancer

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.