

ITA6 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM1828a

Product Information

Application	WB, IHC-P, IF, FC, E
Primary Accession	P23229
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1, κ
Clone Names	129CT49.6.5
Calculated MW	126606

Additional Information

Gene ID	3655
Other Names	Integrin alpha-6, CD49 antigen-like family member F, VLA-6, CD49f, Integrin alpha-6 heavy chain, Integrin alpha-6 light chain, Processed integrin alpha-6, Alpha6p, ITGA6
Target/Specificity	ITA6 recombinant protein is used to produce this monoclonal antibody.
Dilution	WB~1:1000 IHC-P~1:100~500 IF~1:10~50 FC~1:10~50 E~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ITA6 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ITGA6
Function	Integrin alpha-6/beta-1 (ITGA6:ITGB1) is a receptor for laminin on platelets (By similarity). Integrin alpha-6/beta-1 (ITGA6:ITGB1) is present in oocytes and is involved in sperm-egg fusion (By similarity). Integrin alpha-6/beta-4 (ITGA6:ITGB4) is a receptor for laminin in epithelial cells and it plays a critical structural role in the hemidesmosome (By similarity). ITGA6:ITGB4 binds to

NRG1 (via EGF domain) and this binding is essential for NRG1-ERBB signaling (PubMed:[20682778](#)). ITGA6:ITGB4 binds to IGF1 and this binding is essential for IGF1 signaling (PubMed:[22351760](#)). ITGA6:ITGB4 binds to IGF2 and this binding is essential for IGF2 signaling (PubMed:[28873464](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell membrane; Lipid-anchor

Tissue Location

Integrin alpha-6/beta-4 is predominantly expressed by epithelia. Isoforms containing segment X1 are ubiquitously expressed. Isoforms containing segment X1X2 are expressed in heart, kidney, placenta, colon, duodenum, myoblasts and myotubes, and in a limited number of cell lines; they are always coexpressed with the ubiquitous isoform containing segment X1. In some tissues (e.g Salivary gland), isoforms containing cytoplasmic segment A and isoforms containing segment B are detected while in others, only isoforms containing one cytoplasmic segment are found (segment A in epidermis and segment B in kidney). Processed integrin alpha-6: Expressed at low levels in normal prostate tissue with elevated levels in prostate cancer tissue (at protein level) (PubMed:15023541)


Background

The ITGA6 protein product is the integrin alpha chain alpha 6. Integrins are integral cell-surface proteins composed of an alpha chain and a beta chain. A given chain may combine with multiple partners resulting in different integrins. For example, alpha 6 may combine with beta 4 in the integrin referred to as TSP180, or with beta 1 in the integrin VLA-6. Integrins are known to participate in cell adhesion as well as cell-surface mediated signalling. Two transcript variants encoding different isoforms have been found for this gene.


References

Evaluation of candidate stromal epithelial cross-talk genes identifies association between risk of serous ovarian cancer and TERT, a cancer susceptibility hot-spot. Johnatty SE, et al. PLoS Genet, 2010 Jul 8. PMID 20628624. Prostate cancer risk-associated variants reported from genome-wide association studies: Meta-analysis and their contribution to genetic Variation. Kim ST, et al. Prostate, 2010 Jun 16. PMID 20564319. alpha6-integrin subunit plays a major role in the proangiogenic properties of endothelial progenitor cells. Bouvard C, et al. Arterioscler Thromb Vasc Biol, 2010 Aug. PMID 20508204. Common polymorphisms in ITGA2, PON1 and THBS2 are associated with coronary atherosclerosis in a candidate gene association study of the Chinese Han population. Wang Y, et al. J Hum Genet, 2010 Aug. PMID 20485444. Usage of heparan sulfate, integrins, and FAK in HPV16 infection. Abban CY, et al. Virology, 2010 Jul 20. PMID 20441998.


Images




Western blot analysis of anti-ITA6 (Cat. #AM1828a) in HepG2 cell line lysates (35µg/lane). ITA6 (arrow) was detected using the purified Mab.(8µg/ml)



ITA6 Monoclonal Antibody (Cat. #AM1828a) immunohistochemistry analysis in formalin fixed and paraffin embedded human skin carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ITA6 Monoclonal Antibody for immunohistochemistry. Clinical relevance has not been evaluated.



Confocal immunofluorescent analysis of ITA6 Antibody (Cat#AM1828a) with HepG2 cell followed by Alexa Fluor® 488-conjugated goat anti-mouse IgG (green). DAPI was used to stain the cell nuclear (blue).



ITA6 Monoclonal Antibody (Cat. #AM1828a) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). PE-conjugated goat-anti-mouse secondary antibodies were used for the analysis.

Citations

- [Nanoparticle biointerfacing by platelet membrane cloaking.](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.