

LXN Antibody

Mouse Monoclonal Antibody (Mab) Catalog # AM1879b

Product Information

ApplicationWB, EPrimary AccessionQ9BS40Other AccessionNP_064554.3ReactivityHuman, Mouse

HostMouseClonalityMonoclonalIsotypeIgG1,KClone Names199CT6.1.2Calculated MW25750

Additional Information

Gene ID 56925

Other Names Latexin, Endogenous carboxypeptidase inhibitor, ECI, Protein MUM, Tissue

carboxypeptidase inhibitor, TCI, LXN

Target/Specificity This LXN monoclonal antibody is generated from mouse immunized with LXN

recombinant protein.

Dilution WB~~1:100~500 E~~Use at an assay dependent concentration.

Format Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein G column, followed by dialysis

against PBS.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions LXN Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name LXN

Function Hardly reversible, non-competitive, and potent inhibitor of CPA1, CPA2 and

CPA4. May play a role in inflammation.

Cellular Location Cytoplasm.

Highly expressed in heart, prostate, ovary, kidney, pancreas, and colon, moderate or low in other tissues including brain


Background

This gene encodes the only known protein inhibitor of zinc-dependent metallocarboxypeptidases.


References

Lamesch, P., et al. Genomics 89(3):307-315(2007)
Pallares, I., et al. Proc. Natl. Acad. Sci. U.S.A. 102(11):3978-3983(2005)
Liu, Q., et al. Mol. Biol. Rep. 27(4):241-246(2000)

Images



Anti-LXN Antibody at 1:1000 dilution + MCF-7 whole cell lysate Lysates/proteins at 20 μg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-LXN Antibody at 1:250 dilution + mouse brain lysate Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 25750 Da Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.