

# BMI1 Antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AM1930b

## Product Information

---

<b>Application</b>	WB, IF, E
<b>Primary Accession</b>	<a href="#">P35226</a>
<b>Other Accession</b>	<a href="#">NP_005171.4</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1,k
<b>Clone Names</b>	282CT3.7.6
<b>Calculated MW</b>	36949

## Additional Information

---

<b>Gene ID</b>	100532731;648
<b>Other Names</b>	Polycomb complex protein BMI-1, Polycomb group RING finger protein 4, RING finger protein 51, BMI1, PCGF4, RNF51
<b>Target/Specificity</b>	This BMI1 monoclonal antibody is generated from mouse immunized with BMI1 recombinant protein.
<b>Dilution</b>	WB~~1:500~1000 IF~~1:10~50 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	BMI1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	BMI1
<b>Synonyms</b>	PCGF4, RNF51
<b>Function</b>	Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1

complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:[15386022](#), PubMed:[16359901](#), PubMed:[16714294](#), PubMed:[21772249](#), PubMed:[25355358](#), PubMed:[26151332](#), PubMed:[27827373](#)). The complex composed of RNF2, UB2D3 and BMI1 binds nucleosomes, and has activity only with nucleosomal histone H2A (PubMed:[21772249](#), PubMed:[25355358](#)). In the PRC1-like complex, regulates the E3 ubiquitin-protein ligase activity of RNF2/RING2 (PubMed:[15386022](#), PubMed:[21772249](#), PubMed:[26151332](#)).

## Cellular Location

Nucleus. Cytoplasm


## Background

Component of the Polycomb group (PcG) multiprotein PRC1 complex, a complex required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. In the PRC1 complex, it is required to stimulate the E3 ubiquitin-protein ligase activity of RNF2/RING2.


## References

Ismail, I.H., et al. J. Cell Biol. 191(1):45-60(2010)  
 Yang, M.H., et al. Nat. Cell Biol. 12(10):982-992(2010)  
 Kikuchi, J., et al. Cancer 116(12):3015-3024(2010)  
 Honig, A., et al. Anticancer Res. 30(5):1559-1564(2010)  
 Venkataraman, S., et al. PLoS ONE 5 (6), E10748 (2010) :


## Images



All lanes : Anti-BMI1 Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



BMI1 Antibody (Cat. #AM1930b) western blot analysis in K562 cell line lysates (35µg/lane). This demonstrates the BMI1 antibody detected the BMI1 protein (arrow).



Confocal immunofluorescent analysis of BMI1 Antibody (Cat#AM1930b) with NCI-H460 cell followed by Alexa Fluor® 488-conjugated goat anti-mouse IgG (green). Actin filaments have been labeled with Alexa Fluor® 555 phalloidin (red). DAPI was used to stain the cell nuclear (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.