

# UCHL1 Antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AM1959B

## Product Information

---

<b>Application</b>	WB, E
<b>Primary Accession</b>	<a href="#">P09936</a>
<b>Other Accession</b>	<a href="#">NP_004172.2</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1,k
<b>Clone Names</b>	346CT2.2.1
<b>Calculated MW</b>	24824

## Additional Information

---

<b>Gene ID</b>	7345
<b>Other Names</b>	Ubiquitin carboxyl-terminal hydrolase isozyme L1, UCH-L1, 6---, Neuron cytoplasmic protein 95, PGP 95, PGP95, Ubiquitin thioesterase L1, UCHL1
<b>Target/Specificity</b>	This UCHL1 monoclonal antibody is generated from mouse immunized with UCHL1 recombinant protein.
<b>Dilution</b>	WB~~1:120~1000 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	UCHL1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	UCHL1
<b>Function</b>	Deubiquitinase that plays a role in the regulation of several processes such as maintenance of synaptic function, cardiac function, inflammatory response or osteoclastogenesis (PubMed: <a href="#">22212137</a> , PubMed: <a href="#">23359680</a> ). Abrogates the ubiquitination of multiple proteins including WWTR1/TAZ, EGFR, HIF1A and beta-site amyloid precursor protein cleaving enzyme 1/BACE1

(PubMed:[22212137](#), PubMed:[25615526](#)). In addition, recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin to maintain a stable pool of monoubiquitin that is a key requirement for the ubiquitin-proteasome and the autophagy- lysosome pathways (PubMed:[12408865](#), PubMed:[8639624](#), PubMed:[9774100](#)). Regulates amyloid precursor protein/APP processing by promoting BACE1 degradation resulting in decreased amyloid beta production (PubMed:[22212137](#)). Plays a role in the immune response by regulating the ability of MHC I molecules to reach cross-presentation compartments competent for generating Ag-MHC I complexes (By similarity). Mediates the 'Lys-48'-linked deubiquitination of the transcriptional coactivator WWTR1/TAZ leading to its stabilization and inhibition of osteoclastogenesis (By similarity). Deubiquitinates and stabilizes epidermal growth factor receptor EGFR to prevent its degradation and to activate its downstream mediators (By similarity). Modulates oxidative activity in skeletal muscle by regulating key mitochondrial oxidative proteins (By similarity). Enhances the activity of hypoxia-inducible factor 1-alpha/HIF1A by abrogating its VHL E3 ligase-mediated ubiquitination and consequently inhibiting its degradation (PubMed:[25615526](#)).

#### Cellular Location

Cytoplasm. Endoplasmic reticulum membrane; Lipid- anchor. Note=About 30% of total UCHL1 is associated with membranes in brain. Localizes near and/or within mitochondria to potentially interact with mitochondrial proteins {ECO:0000250|UniProtKB:Q9R0P9}

#### Tissue Location

Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.

## Background


The protein encoded by this gene belongs to the peptidase C12 family. This enzyme is a thiol protease that hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. This gene is specifically expressed in the neurons and in cells of the diffuse neuroendocrine system. Mutations in this gene may be associated with Parkinson disease.

## References

- Martins-de-Souza, D., et al. J Psychiatr Res 44(14):989-991(2010)  
 Hussain, S., et al. Leukemia 24(9):1641-1655(2010)  
 Ma, Y., et al. J. Cell. Biochem. 110(6):1512-1519(2010)  
 Wu, Y.R., et al. Clin. Chim. Acta 411 (13-14), 955-958 (2010) :  
 Li, L., et al. Clin. Cancer Res. 16(11):2949-2958(2010)

## Images

All lanes :UCHL1 Antibody at 1:2000 dilution Lane 1:  
 Mouse brain lysate Lane 2: C6 whole cell lysate Lane 3:  
 NCI-H460 whole cell lysate Lysates/proteins at 20 µg per  
 lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase  
 conjugated at 1/10000 dilution. Predicted band size : 25  
 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.