

EIF4B Antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AM1962b

Product Information

Application	WB, E
Primary Accession	P23588
Other Accession	NP_001408.2
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,k
Clone Names	355CT12.6.5
Calculated MW	69151

Additional Information

Gene ID	1975
Other Names	Eukaryotic translation initiation factor 4B, eIF-4B, EIF4B
Target/Specificity	This EIF4B monoclonal antibody is generated from mouse immunized with EIF4B recombinant protein.
Dilution	WB~~ 1:500 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	EIF4B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	EIF4B
Function	Required for the binding of mRNA to ribosomes. Functions in close association with EIF4-F and EIF4-A. Binds near the 5'-terminal cap of mRNA in presence of EIF-4F and ATP. Promotes the ATPase activity and the ATP-dependent RNA unwinding activity of both EIF4-A and EIF4-F.


Background

Required for the binding of mRNA to ribosomes. Functions in close association with EIF4-F and EIF4-A. Binds near the 5'-terminal cap of mRNA in presence of EIF-4F and ATP. Promotes the ATPase activity and the ATP-dependent RNA unwinding activity of both EIF4-A and EIF4-F.


References

Mokas, S., et al. Mol. Biol. Cell 20(11):2673-2683(2009)
Rozovsky, N., et al. RNA 14(10):2136-2148(2008)
Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Olsen, J.V., et al. Cell 127(3):635-648(2006)

Images



All lanes: Anti-EIF4B Antibody at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Observed band size: 80 KDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes: Anti-EIF4B Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: 293 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Observed band size: 80 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.