

# GAD2 Antibody (Ascites)

Mouse Monoclonal Antibody (Mab)

Catalog # AM2049a

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">Q05329</a>
Other Accession	<a href="#">Q05683</a> , <a href="#">P48321</a> , <a href="#">P48320</a> , <a href="#">NP_000809.1</a>
Reactivity	Human, Rat
Predicted	Mouse, Pig
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Names	458CT17.2.4
Calculated MW	65411
Antigen Region	109-138

## Additional Information

---

Gene ID	2572
Other Names	Glutamate decarboxylase 2, 65 kDa glutamic acid decarboxylase, GAD-65, Glutamate decarboxylase 65 kDa isoform, GAD2, GAD65
Target/Specificity	This GAD2 antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 109-138 amino acids from human GAD2.
Dilution	WB~~1:100~1600 E~~Use at an assay dependent concentration.
Format	Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GAD2 Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	GAD2 ( <a href="#">HGNC:4093</a> )
Synonyms	GAD65
Function	Catalyzes the production of GABA.

## Cellular Location

Cytoplasm, cytosol. Cytoplasmic vesicle. Presynaptic cell membrane; Lipid-anchor. Golgi apparatus membrane; Peripheral membrane protein; Cytoplasmic side. Note=Associated to cytoplasmic vesicles In neurons, cytosolic leaflet of Golgi membranes and presynaptic clusters

## Background

---

This gene encodes one of several forms of glutamic acid decarboxylase, identified as a major autoantigen in insulin-dependent diabetes. The enzyme encoded is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantibody and an autoreactive T cell target in insulin-dependent diabetes. This gene may also play a role in the stiff man syndrome. Alternative splicing results in multiple transcript variants that encode the same protein.


## References

---

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)  
Jia, P., et al. Schizophr. Res. 122 (1-3), 38-42 (2010) :  
Ruano, G., et al. Pharmacogenomics 11(7):959-971(2010)  
Pinheiro, A.P., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (5), 1070-1080 (2010) :  
Jugessur, A., et al. PLoS ONE 5 (7), E11493 (2010) :

## Images

---



GAD2 Antibody (Cat. #AM2049a) western blot analysis in rat brain tissue lysates (35µg/lane). This demonstrates the GAD2 antibody detected the GAD2 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.