10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

B2M Antibody (Ascites)

Mouse Monoclonal Antibody (Mab)
Catalog # AM2052a

Product Information

Application	WB, E
Primary Accession	P61769
Other Accession	NP_004039.1
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Names	467CT12.3.1
Calculated MW	13715
Antigen Region	10-39

Additional Information

Gene ID	567
Other Names	Beta-2-microglobulin, Beta-2-microglobulin form pI 53, B2M
Target/Specificity	This B2M antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 10-39 amino acids from human B2M.
Dilution	WB~~1:1000~16000 E~~Use at an assay dependent concentration.
Format	Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	B2M Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	B2M (HGNC:914)
Function	Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system. Exogenously applied M.tuberculosis EsxA or EsxA-EsxB (or EsxA expressed in host) binds B2M and decreases its export to the cell surface (total protein levels do not change), probably leading to defects in class I antigen presentation

(PubMed:25356553).

Cellular Location Secreted. Cell surface. Note=Detected in serum and urine (PubMed:1336137, PubMed:7554280). {ECO:0000269|PubMed:7554280, ECO:0000269|Ref.6}


Background

This gene encodes a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells. The protein has a predominantly beta-pleated sheet structure that can form amyloid fibrils in some pathological conditions. A mutation in this gene has been shown to result in hypercatabolic hypoproteinemia.

References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Rennella, E., et al. J. Mol. Biol. 401(2):286-297(2010)
Debelouchina, G.T., et al. J. Am. Chem. Soc. 132(30):10414-10423(2010)
Mumtaz, A., et al. Saudi J Kidney Dis Transpl 21(4):701-706(2010)
Guo, H.C., et al. Nature 360(6402):364-366(1992)

Images



B2M Antibody (Cat. #AM2052a) western blot analysis in CEM cell line lysates (35µg/lane).This demonstrates the B2M antibody detected the B2M protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.