

B2M Antibody

Mouse Monoclonal Antibody (Mab)

Catalog # AM2052b

Product Information

Application	WB, E
Primary Accession	P61769
Other Accession	NP_004039.1
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Names	467CT12.3.1
Calculated MW	13715
Antigen Region	10-39

Additional Information

Gene ID	567
Other Names	Beta-2-microglobulin, Beta-2-microglobulin form pI 53, B2M
Target/Specificity	This B2M antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 10-39 amino acids from human B2M.
Dilution	WB~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	B2M Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	B2M (HGNC:914)
Function	Component of the class I major histocompatibility complex (MHC). Involved in the presentation of peptide antigens to the immune system. Exogenously applied M.tuberculosis EsxA or EsxA-EsxB (or EsxA expressed in host) binds B2M and decreases its export to the cell surface (total protein levels do not change), probably leading to defects in class I antigen presentation

(PubMed:[25356553](#)).

Cellular Location

Secreted. Cell surface. Note=Detected in serum and urine (PubMed:1336137, PubMed:7554280). {ECO:0000269 | PubMed:7554280, ECO:0000269 | Ref.6}


Background

This gene encodes a serum protein found in association with the major histocompatibility complex (MHC) class I heavy chain on the surface of nearly all nucleated cells. The protein has a predominantly beta-pleated sheet structure that can form amyloid fibrils in some pathological conditions. A mutation in this gene has been shown to result in hypercatabolic hypoproteinemia.

References

- Bailey, S.D., et al. *Diabetes Care* 33(10):2250-2253(2010)
Rennella, E., et al. *J. Mol. Biol.* 401(2):286-297(2010)
Debelouchina, G.T., et al. *J. Am. Chem. Soc.* 132(30):10414-10423(2010)
Mumtaz, A., et al. *Saudi J Kidney Dis Transpl* 21(4):701-706(2010)
Guo, H.C., et al. *Nature* 360(6402):364-366(1992)

Images



B2M Antibody (N-term) (MS2134) western blot analysis in Jurkat, HeLa cell line lysates (35µg/lane). This demonstrates the B2M antibody detected the B2M protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.