

ACE Antibody (Ascites)

Mouse Monoclonal Antibody (Mab)

Catalog # AM2111a

Product Information

Application	WB, E
Primary Accession	P12821
Other Accession	NP_000780
Reactivity	Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Clone Names	536CT24.1.3
Calculated MW	149715
Antigen Region	1274-1306

Additional Information

Gene ID	1636
Other Names	Angiotensin-converting enzyme, ACE, 321-, Dipeptidyl carboxypeptidase I, Kininase II, CD143, Angiotensin-converting enzyme, soluble form, ACE, DCP, DCP1
Target/Specificity	This ACE antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 1274-1306 amino acids from human ACE.
Dilution	WB~~1:500~16000 E~~Use at an assay dependent concentration.
Format	Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ACE Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ACE {ECO:0000303 PubMed:2849100, ECO:0000312 HGNC:HGNC:2707}
Function	Dipeptidyl carboxypeptidase that removes dipeptides from the C-terminus of a variety of circulating hormones, such as angiotensin I, bradykinin or enkephalins, thereby playing a key role in the regulation of blood pressure, electrolyte homeostasis or synaptic plasticity (PubMed: 15615692 ,

PubMed:[20826823](#), PubMed:[2558109](#), PubMed:[4322742](#), PubMed:[7523412](#), PubMed:[7683654](#)). Composed of two similar catalytic domains, each possessing a functional active site, with different selectivity for substrates (PubMed:[10913258](#), PubMed:[1320019](#), PubMed:[1851160](#), PubMed:[19773553](#), PubMed:[7683654](#), PubMed:[7876104](#)). Plays a major role in the angiotensin-renin system that regulates blood pressure and sodium retention by the kidney by converting angiotensin I to angiotensin II, resulting in an increase of the vasoconstrictor activity of angiotensin (PubMed:[11432860](#), PubMed:[1851160](#), PubMed:[19773553](#), PubMed:[23056909](#), PubMed:[4322742](#)). Also able to inactivate bradykinin, a potent vasodilator, and therefore enhance the blood pressure response (PubMed:[15615692](#), PubMed:[2558109](#), PubMed:[4322742](#), PubMed:[6055465](#), PubMed:[6270633](#), PubMed:[7683654](#)). Acts as a regulator of synaptic transmission by mediating cleavage of neuropeptide hormones, such as substance P, neurotensin or enkephalins (PubMed:[15615692](#), PubMed:[6208535](#), PubMed:[6270633](#), PubMed:[656131](#)). Catalyzes degradation of different enkephalin neuropeptides (Met-enkephalin, Leu-enkephalin, Met-enkephalin-Arg-Phe and possibly Met-enkephalin-Arg-Gly-Leu) (PubMed:[2982830](#), PubMed:[6270633](#), PubMed:[656131](#)). Acts as a regulator of synaptic plasticity in the nucleus accumbens of the brain by mediating cleavage of Met-enkephalin- Arg-Phe, a strong ligand of Mu-type opioid receptor OPRM1, into Met- enkephalin (By similarity). Met-enkephalin-Arg-Phe cleavage by ACE decreases activation of OPRM1, leading to long-term synaptic potentiation of glutamate release (By similarity). Also acts as a regulator of hematopoietic stem cell differentiation by mediating degradation of hemoregulatory peptide N-acetyl-SDKP (AcSDKP) (PubMed:[26403559](#), PubMed:[7876104](#), PubMed:[8257427](#), PubMed:[8609242](#)). Acts as a regulator of cannabinoid signaling pathway by mediating degradation of hemopressin, an antagonist peptide of the cannabinoid receptor CNR1 (PubMed:[18077343](#)). Involved in amyloid-beta metabolism by catalyzing degradation of Amyloid-beta protein 40 and Amyloid-beta protein 42 peptides, thereby preventing plaque formation (PubMed:[11604391](#), PubMed:[16154999](#), PubMed:[19773553](#)). Catalyzes cleavage of cholecystokinin (maturation of Cholecystokinin-8 and Cholecystokinin-5) and Gonadoliberin-1 (both maturation and degradation) hormones (PubMed:[10336644](#), PubMed:[2983326](#), PubMed:[7683654](#), PubMed:[9371719](#)). Degradation of hemoregulatory peptide N-acetyl-SDKP (AcSDKP) and amyloid-beta proteins is mediated by the N-terminal catalytic domain, while angiotensin I and cholecystokinin cleavage is mediated by the C-terminal catalytic region (PubMed:[10336644](#), PubMed:[19773553](#), PubMed:[7876104](#)).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cytoplasm {ECO:0000250|UniProtKB:P09470}. Note=Detected in both cell membrane and cytoplasm in neurons. {ECO:0000250|UniProtKB:P09470} [Isoform Testis-specific]: Cell membrane; Single-pass type I membrane protein. Secreted. Note=The testis-specific isoform can be cleaved before the transmembrane region, releasing a soluble form

Tissue Location

Ubiquitously expressed, with highest levels in lung, kidney, heart, gastrointestinal system and prostate

Background


This gene encodes an enzyme involved in catalyzing the conversion of angiotensin I into a physiologically active peptide angiotensin II. Angiotensin II is a potent vasopressor and aldosterone-stimulating peptide that controls blood pressure and fluid-electrolyte balance. This enzyme plays a key role in the renin-angiotensin system. Many studies have associated the presence or absence of a 287 bp Alu repeat element in this gene with the levels of circulating enzyme or cardiovascular pathophysiology. Multiple alternatively spliced transcript variants encoding different isoforms have been identified, and two most abundant spliced variants encode the somatic form and the testicular form, respectively, that are equally

active.

References

Dimitriou, G., et al. *Pediatr. Pulmonol.* 45(12):1233-1239(2010)
Ince, D.A., et al. *Genet Test Mol Biomarkers* 14(5):643-647(2010)
Procopciuc, L.M., et al. *Eur. J. Intern. Med.* 21(5):414-418(2010)
Ash, G.I., et al. *Med Sci Sports Exerc* (2010) In press :
Liu, L.W., et al. *Chin. Med. J.* 123(11):1382-1386(2010)

Images



ACE Antibody (Ascites)(Cat. #AM2111a) western blot analysis in mouse kidney tissue lysates (35µg/lane). This demonstrates the ACE antibody detected the ACE protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.