

UCHL1 Antibody (C-term)

Mouse Monoclonal Antibody (Mab)

Catalog # AM2207b

Product Information

Application	WB, IHC-P, E
Primary Accession	P09936
Other Accession	Q00981 , Q6SEG5 , Q9R0P9 , Q60HC8 , P23356
Reactivity	Human, Rat, Mouse
Predicted	Bovine, Monkey, Mouse, Pig
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Clone Names	696CT2.1.4
Calculated MW	24824

Additional Information

Gene ID	7345
Other Names	Ubiquitin carboxyl-terminal hydrolase isozyme L1, UCH-L1, 6---, Neuron cytoplasmic protein 95, PGP 95, PGP95, Ubiquitin thioesterase L1, UCHL1
Target/Specificity	This UCHL1 Monoclonal antibody is generated from mice immunized with a KLH conjugated synthetic peptide selected from the 185-214 region of human UCHL1.
Dilution	WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	UCHL1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	UCHL1
Function	Deubiquitinase that plays a role in the regulation of several processes such as maintenance of synaptic function, cardiac function, inflammatory response or osteoclastogenesis (PubMed: 22212137 , PubMed: 23359680). Abrogates the

ubiquitination of multiple proteins including WWTR1/TAZ, EGFR, HIF1A and beta-site amyloid precursor protein cleaving enzyme 1/BACE1 (PubMed:[22212137](#), PubMed:[25615526](#)). In addition, recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin to maintain a stable pool of monoubiquitin that is a key requirement for the ubiquitin-proteasome and the autophagy- lysosome pathways (PubMed:[12408865](#), PubMed:[8639624](#), PubMed:[9774100](#)). Regulates amyloid precursor protein/APP processing by promoting BACE1 degradation resulting in decreased amyloid beta production (PubMed:[22212137](#)). Plays a role in the immune response by regulating the ability of MHC I molecules to reach cross-presentation compartments competent for generating Ag-MHC I complexes (By similarity). Mediates the 'Lys-48'-linked deubiquitination of the transcriptional coactivator WWTR1/TAZ leading to its stabilization and inhibition of osteoclastogenesis (By similarity). Deubiquitinates and stabilizes epidermal growth factor receptor EGFR to prevent its degradation and to activate its downstream mediators (By similarity). Modulates oxidative activity in skeletal muscle by regulating key mitochondrial oxidative proteins (By similarity). Enhances the activity of hypoxia-inducible factor 1-alpha/HIF1A by abrogating its VHL E3 ligase-mediated ubiquitination and consequently inhibiting its degradation (PubMed:[25615526](#)).

Cellular Location

Cytoplasm. Endoplasmic reticulum membrane; Lipid- anchor. Note=About 30% of total UCHL1 is associated with membranes in brain. Localizes near and/or within mitochondria to potentially interact with mitochondrial proteins {ECO:0000250|UniProtKB:Q9R0P9}

Tissue Location

Found in neuronal cell bodies and processes throughout the neocortex (at protein level). Expressed in neurons and cells of the diffuse neuroendocrine system and their tumors. Weakly expressed in ovary. Down-regulated in brains from Parkinson disease and Alzheimer disease patients.

Background



Ubiquitin-protein hydrolase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. This enzyme is a thiol protease that recognizes and hydrolyzes a peptide bond at the C-terminal glycine of ubiquitin. Also binds to free monoubiquitin and may prevent its degradation in lysosomes. The homodimer may have ATP-independent ubiquitin ligase activity.

References


- Boudreaux D.A., et al. Proc. Natl. Acad. Sci. U.S.A. 107:9117-9122(2010).
 Hillier L.W., et al. Nature 434:724-731(2005).
 Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
 Day I.N.M., et al. Biochem. J. 268:521-524(1990).
 Choi J., et al. J. Biol. Chem. 279:13256-13264(2004).

Images

Immunohistochemical analysis of paraffin-embedded H. brain section using UCHL1 Antibody (C-term)(Cat#AM2207B). AM2207B was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. astrogloma section using UCHL1 Antibody (C-term)(Cat#AM2207B). AM2207B was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



UCHL1 Antibody (C-term)(Cat. #AM2207b) western blot analysis in 293T,NCI-H1299,rat PC-12 cell line lysates (35µg/lane).This demonstrates the UCHL1 antibody detected the UCHL1 protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.