

# LCK Antibody(Ascites)

Mouse Monoclonal Antibody (Mab)  
Catalog # AM2213a

## Product Information

---

|                          |                        |
|--------------------------|------------------------|
| <b>Application</b>       | WB, E                  |
| <b>Primary Accession</b> | <a href="#">P06239</a> |
| <b>Reactivity</b>        | Human                  |
| <b>Host</b>              | Mouse                  |
| <b>Clonality</b>         | Monoclonal             |
| <b>Isotype</b>           | IgG1                   |
| <b>Clone Names</b>       | 845CT3.5.2             |
| <b>Calculated MW</b>     | 58001                  |

## Additional Information

---

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 3932   |
| <b>Other Names</b>        | Tyrosine-protein kinase Lck, Leukocyte C-terminal Src kinase, LSK, Lymphocyte cell-specific protein-tyrosine kinase, Protein YT16, Proto-oncogene Lck, T cell-specific protein-tyrosine kinase, p56-LCK, LCK |
| <b>Target/Specificity</b> | Purified His-tagged LCK protein was used to produced this monoclonal antibody.   |
| <b>Dilution</b>           | WB~~1:5000 E~~Use at an assay dependent concentration.   |
| <b>Format</b>             | Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.   |
| <b>Storage</b>            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.  |
| <b>Precautions</b>        | LCK Antibody(Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.  |

## Protein Information

---

|                 |  |
|-----------------|--|
| <b>Name</b>     | LCK  |
| <b>Function</b> | Non-receptor tyrosine-protein kinase that plays an essential role in the selection and maturation of developing T-cells in the thymus and in the function of mature T-cells (PubMed: <a href="#">2470098</a> ). Plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways (PubMed: <a href="#">2470098</a> ). Constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors (PubMed: <a href="#">2470098</a> ). Association of the TCR with a peptide |

antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, thereby recruiting the associated LCK protein to the vicinity of the TCR-CD3 complex (PubMed:[2470098](#)). LCK then phosphorylates tyrosine residues within the immunoreceptor tyrosine-based activation motifs (ITAM) of the cytoplasmic tails of the TCR-gamma chains and CD3 subunits, initiating the TCR-CD3 signaling pathway (PubMed:[2470098](#), PubMed:[40592325](#)). Once stimulated, the TCR recruits the tyrosine kinase ZAP70, that becomes phosphorylated and activated by LCK. Following this, a large number of signaling molecules are recruited, ultimately leading to lymphokine production. LCK also contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, which leads to hyperphosphorylation and activation of LCK. Also plays a role in the IL2 receptor-linked signaling pathway that controls the T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Phosphorylates other substrates including RUNX3, PTK2B/PYK2, the microtubule-associated protein MAPT, RHOH or TYROBP. Interacts with FYB2 (PubMed:[27335501](#)).

#### **Cellular Location**

Cell membrane; Lipid-anchor; Cytoplasmic side Cytoplasm, cytosol.  
Note=Present in lipid rafts in an inactive form.

#### **Tissue Location**

Expressed specifically in lymphoid cells.

## **Background**

---

Non-receptor tyrosine-protein kinase that plays an essential role in the selection and maturation of developing T-cells in the thymus and in the function of mature T-cells. Plays a key role in T-cell antigen receptor (TCR)-linked signal transduction pathways. Constitutively associated with the cytoplasmic portions of the CD4 and CD8 surface receptors. Association of the TCR with a peptide antigen-bound MHC complex facilitates the interaction of CD4 and CD8 with MHC class II and class I molecules, respectively, thereby recruiting the associated LCK protein to the vicinity of the TCR/CD3 complex. LCK then phosphorylates tyrosines residues within the immunoreceptor tyrosine-based activation motifs (ITAM) of the cytoplasmic tails of the TCR-gamma chains and CD3 subunits, initiating the TCR/CD3 signaling pathway. Once stimulated, the TCR recruits the tyrosine kinase ZAP70, that becomes phosphorylated and activated by LCK. Following this, a large number of signaling molecules are recruited, ultimately leading to lymphokine production. LCK also contributes to signaling by other receptor molecules. Associates directly with the cytoplasmic tail of CD2, which leads to hyperphosphorylation and activation of LCK. Also plays a role in the IL2 receptor-linked signaling pathway that controls the T-cell proliferative response. Binding of IL2 to its receptor results in increased activity of LCK. Is expressed at all stages of thymocyte development and is required for the regulation of maturation events that are governed by both pre-TCR and mature alpha beta TCR. Phosphorylates other substrates including RUNX3, PTK2B/PYK2, the microtubule-associated protein MAPT, RHOH or TYROBP.

## **References**


---

Yamaguchi H., et al. *Nature* 384:484-489(1996).  
Koga Y., et al. *Eur. J. Immunol.* 16:1643-1646(1986).  
Perlmutter R.M., et al. *J. Cell. Biochem.* 38:117-126(1988).  
Rouer E., et al. *Gene* 84:105-113(1989).  
Wright D.D., et al. *Mol. Cell. Biol.* 14:2429-2437(1994).

## **Images**

---

LCK Antibody(Cat. #AM2213a) western blot analysis in



HL-60,Jurkat,NCI-H292 cell line lysates (35 $\mu$ g/lane).This demonstrates the LCK antibody detected the LCK protein (arrow).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.