

# SUMO2 Antibody (C-term)

Mouse Monoclonal Antibody (Mab)

Catalog # AM2225b

## Product Information

---

|                   |  |
|-------------------|--|
| Application       | WB, IF, E  |
| Primary Accession | <a href="#">P61956</a>   |
| Other Accession   | <a href="#">P61959</a> , <a href="#">P61958</a> , <a href="#">P61957</a> , <a href="#">Q2PFW2</a> , <a href="#">Q6LDZ8</a> , <a href="#">Q5ZJM9</a> , <a href="#">P61955</a> |
| Reactivity        | Human, Rat   |
| Predicted         | Bovine, Chicken, Hamster, Monkey, Mouse, Pig   |
| Host              | Mouse  |
| Clonality         | Monoclonal   |
| Isotype           | IgG2b  |
| Clone Names       | 973CT8.1.1   |
| Calculated MW     | 10871  |

## Additional Information

---

|                    |   |
|--------------------|---|
| Gene ID            | 6613  |
| Other Names        | Small ubiquitin-related modifier 2, SUMO-2, HSMT3, SMT3 homolog 2 {ECO:0000312   HGNC:HGNC:11125}, SUMO-3, Sentrin-2, Ubiquitin-like protein SMT3B, Smt3B, SUMO2 ( <a href="#">HGNC:11125</a> ) |
| Target/Specificity | Purified His-tagged SUMO2 protein was used to produced this monoclonal antibody.  |
| Dilution           | WB~~1:1000 IF~~1:25 E~~Use at an assay dependent concentration.   |
| Format             | Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.                             |
| Storage            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.   |
| Precautions        | SUMO2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.   |

## Protein Information

---

|          |   |
|----------|---|
| Name     | SUMO2 ( <a href="#">HGNC:11125</a> )  |
| Function | Ubiquitin-like protein that can be covalently attached to proteins as a monomer or as a lysine-linked polymer. Covalent attachment via an isopeptide bond to its substrates requires prior activation by the E1 complex |

SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by an E3 ligase such as PIAS1-4, RANBP2, CBX4 or ZNF451 (PubMed:[26524494](#)). This post-translational modification on lysine residues of proteins plays a crucial role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Polymeric SUMO2 chains are also susceptible to polyubiquitination which functions as a signal for proteasomal degradation of modified proteins (PubMed:[18408734](#), PubMed:[18538659](#), PubMed:[21965678](#), PubMed:[9556629](#)). Plays a role in the regulation of sumoylation status of SETX (PubMed:[24105744](#)).

**Cellular Location** Nucleus. Nucleus, PML body.

**Tissue Location** Broadly expressed..

## Background

---

Ubiquitin-like protein that can be covalently attached to proteins as a monomer or as a lysine-linked polymer. Covalent attachment via an isopeptide bond to its substrates requires prior activation by the E1 complex SAE1-SAE2 and linkage to the E2 enzyme UBE2I, and can be promoted by an E3 ligase such as PIAS1-4, RANBP2 or CBX4. This post-translational modification on lysine residues of proteins plays a crucial role in a number of cellular processes such as nuclear transport, DNA replication and repair, mitosis and signal transduction. Polymeric SUMO2 chains are also susceptible to polyubiquitination which functions as a signal for proteasomal degradation of modified proteins.


## References

---

- Reverter D., et al. Structure 12:1519-1531(2004).  
Xu Z., et al. Biochem. J. 386:325-330(2005).  
Mannen H., et al. Biochem. Biophys. Res. Commun. 222:178-180(1996).  
Lapenta V., et al. Genomics 40:362-367(1997).  
Ota T., et al. Nat. Genet. 36:40-45(2004).


## Images

---



Fluorescent image of HeLa cells stained with SUMO2 Antibody (C-term)(Cat#AM2225B). AM2225B was diluted at 1:25 dilution. An Alexa Fluor® 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).

SUMO2 Antibody (C-term)(Cat. #AM2225b) western blot analysis in CEM,293, rat C6 cell line lysates (35µg/lane). This demonstrates the SUMO2 antibody detected the SUMO2 protein (arrow).



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.