10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

RAD51 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8421b

Product Information

Application	WB, E
Primary Accession	Q06609
Reactivity	Human, Mouse, Rat
Predicted	Bovine, Canine, Rabbit, Chicken
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,κ
Clone Names	1281CT886.273.179.159
Calculated MW	36966
Antigen Region	1-250

Additional Information

Gene ID	5888
Other Names	DNA repair protein RAD51 homolog 1, HsRAD51, hRAD51, RAD51 homolog A, RAD51, RAD51A, RECA
Target/Specificity	This RAD51 antibody is generated from a mouse immunized with a recombination protein from human.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RAD51 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RAD51 {ECO:0000303|PubMed:8358431, ECO:0000312|HGNC:HGNC:9817}
Function	Homologous DNA recombinase that catalyzes strand exchange, a key step in DNA repair through homologous recombination (HR) (PubMed:12205100, PubMed:12442171, PubMed:15937124, PubMed:17515904, PubMed:17515903, PubMed:18417535, PubMed:19303847,

PubMed:20231364, PubMed:20348101, PubMed:22325354, PubMed:23509288, PubMed:23754376, PubMed:24141787, PubMed:26681308, PubMed:27941124, PubMed:28575658, PubMed:32640219, PubMed:37499663, PubMed:38509361, PubMed:7988572). Binds to single-stranded DNA (ssDNA) in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed:12205100, PubMed:15937124, PubMed:17515904, PubMed:17515903, PubMed:18417535, PubMed:19303847, PubMed:15226506, PubMed:20231364, PubMed:20348101, PubMed:23509288, PubMed:23754376, PubMed:26681308, PubMed:28575658, PubMed:37499663, PubMed:39636933, PubMed:41166468, PubMed:38509361, PubMed:7988572). Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template (PubMed:12205100, PubMed:18417535, PubMed:19303847, PubMed:20231364, PubMed:20348101, PubMed:23509288, PubMed:23754376, PubMed:26681308, PubMed:28575658, PubMed:38459011). RAD51 targeting to ssDNA promotes removal of replication protein-A (RPA) from ssDNA and stabilization of RAD51-ssDNA filaments by blocking ATP hydrolysis (PubMed:20729859). Nucleosomal DNA is peeled from the histone surface by the RAD51 filament (PubMed:38509361). Recruited to resolve stalled replication forks during replication stress (PubMed:27797818, PubMed:31844045). Also required for homologous recombination during meiosis by acting as an non-catalytic accessory factor for DMC1 recombinase, enabling proper chromosome pairing and crossing over (By similarity). Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3 (PubMed:20413593). Also involved in interstrand cross-link repair (PubMed:26253028).

Cellular Location

Chromosome. Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Mitochondrion matrix. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Colocalizes with RAD51AP1 and RPA2 to multiple nuclear foci upon induction of DNA damage (PubMed:20154705). DNA damage induces an increase in nuclear levels (PubMed:20154705). Together with FIGNL1, redistributed in discrete nuclear DNA damage-induced foci after ionizing radiation (IR) or camptothecin (CPT) treatment (PubMed:23754376). Accumulated at sites of DNA damage in a SPIDR- dependent manner (PubMed:23509288). Recruited at sites of DNA damage in a MCM9-MCM8-dependent manner (PubMed:23401855). Recruited at sites of DNA damage following interaction with TOPBP1 in S-phase (PubMed:26811421). Colocalizes with ERCC5/XPG to nuclear foci in S phase (PubMed:26833090). Recruited to stalled replication forks during replication stress by the TONSL-MMS22L complex, as well as ATAD5 and WDR48 in an ATR-dependent manner (PubMed:27797818, PubMed:31844045). In germ cells, during meiotic prcess, binds single-stranded DNA (ssDNA) away from the break site (By similarity) {ECO:0000250|UniProtKB:Q08297, ECO:0000269|PubMed:20154705, ECO:0000269|PubMed:23401855, ECO:0000269|PubMed:23509288, ECO:0000269|PubMed:23754376, ECO:0000269|PubMed:26811421, ECO:0000269|PubMed:26833090, ECO:0000269|PubMed:27797818, ECO:0000269|PubMed:31844045}

Tissue Location

Highly expressed in testis and thymus, followed by small intestine, placenta, colon, pancreas and ovary (PubMed:18417535) Weakly expressed in breast (PubMed:18417535)


Background

Participates in a common DNA damage response pathway associated with the activation of homologous recombination and double-strand break repair. Binds to single and double-stranded DNA and exhibits DNA-dependent ATPase activity. Underwinds duplex DNA and forms helical nucleoprotein filaments. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3.

References

Shinohara A.,et al.Nat. Genet. 4:239-243(1993).
Yoshimura Y.,et al.Nucleic Acids Res. 21:1665-1665(1993).
Schmutte C.,et al.Cancer Res. 59:4564-4569(1999).
Wang W.W.,et al.Cancer Epidemiol. Biomarkers Prev. 10:955-960(2001).
Park J.Y.,et al.Nucleic Acids Res. 36:3226-3234(2008).

Images



All lanes: Anti-RAD51 at 1:2000 dilution Lane 1: C2C12 whole cell lysate Lane 2: C6 whole cell lysate Lane 3: Mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Observed band size: 35 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.