

RPTOR Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8442b

Product Information

Application	WB, IHC-P, E
Primary Accession	Q8N122
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,k
Clone Names	1411CT316.2.151.34
Calculated MW	149038
Antigen Region	1005-1329

Additional Information

Gene ID	57521
Other Names	Regulatory-associated protein of mTOR, Raptor, p150 target of rapamycin (TOR)-scaffold protein, RPTOR, KIAA1303, RAPTOR
Target/Specificity	This RPTOR antibody is generated from a mouse immunized with recombinant protein.
Dilution	WB~~1:500-1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RPTOR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RPTOR (HGNC:30287)
Function	Component of the mechanistic target of rapamycin complex 1 (mTORC1), an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed: 12150925 , PubMed: 12150926 ,

PubMed:[12747827](#), PubMed:[24403073](#), PubMed:[26588989](#), PubMed:[32561715](#), PubMed:[37541260](#)). In response to nutrients, growth factors or amino acids, mTORC1 is recruited to the lysosome membrane and promotes protein, lipid and nucleotide synthesis by phosphorylating several substrates, such as ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) (PubMed:[12150925](#), PubMed:[12150926](#), PubMed:[12747827](#), PubMed:[24403073](#), PubMed:[26588989](#), PubMed:[37541260](#)). In the same time, it inhibits catabolic pathways by phosphorylating the autophagy initiation components ULK1 and ATG13, as well as transcription factor TFEB, a master regulators of lysosomal biogenesis and autophagy (PubMed:[12150925](#), PubMed:[12150926](#), PubMed:[12747827](#), PubMed:[24403073](#), PubMed:[32561715](#), PubMed:[37541260](#)). The mTORC1 complex is inhibited in response to starvation and amino acid depletion (PubMed:[12150925](#), PubMed:[12150926](#), PubMed:[12747827](#), PubMed:[24403073](#), PubMed:[37541260](#)). Within the mTORC1 complex, RPTOR acts both as a molecular adapter, which (1) mediates recruitment of mTORC1 to lysosomal membranes via interaction with small GTPases Rag (RagA/RRAGA, RagB/RRAGB, RagC/RRAGC and/or RagD/RRAGD), and a (2) substrate-specific adapter, which promotes substrate specificity by binding to TOS motif- containing proteins and direct them towards the active site of the MTOR kinase domain for phosphorylation (PubMed:[12747827](#), PubMed:[24403073](#), PubMed:[26588989](#), PubMed:[37541260](#)). mTORC1 complex regulates many cellular processes, such as odontoblast and osteoclast differentiation or neuronal transmission (By similarity). mTORC1 complex in excitatory neuronal transmission is required for the prosocial behavior induced by the psychoactive substance lysergic acid diethylamide (LSD) (By similarity).

Cellular Location

Lysosome membrane. Cytoplasm Cytoplasmic granule. Note=Targeting to lysosomes depends on amino acid availability: recruited to lysosome membranes via interaction with GTP-bound form of RagA/RRAGA (or RagB/RRAGB) in complex with the GDP-bound form of RagC/RRAGC (or RagD/RRAGD), promoting recruitment of mTORC1 to the lysosomes (PubMed:31601708, PubMed:31601764). In arsenite-stressed cells, accumulates in stress granules when associated with SPAG5 and association with lysosomes is drastically decreased (PubMed:23953116).

Tissue Location

Highly expressed in skeletal muscle, and in a lesser extent in brain, lung, small intestine, kidney and placenta

Background


Involved in the control of the mammalian target of rapamycin complex 1 (mTORC1) activity which regulates cell growth and survival, and autophagy in response to nutrient and hormonal signals; functions as a scaffold for recruiting mTORC1 substrates. mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated mTORC1 activation involves a AKT1- mediated phosphorylation of TSC1-TSC2, which leads to the activation of the RHEB GTPase that potently activates the protein kinase activity of mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation initiation factor 4E (eIF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation. Involved in ciliogenesis.

References


Kim D.-H.,et al.Cell 110:163-175(2002).
Hara K.,et al.Cell 110:177-189(2002).

Zody M.C.,et al.Nature 440:1045-1049(2006).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Nagase T.,et al.DNA Res. 7:65-73(2000).

Images



Immunohistochemical analysis of paraffin-embedded H. skeletal muscle section using RPTOR Antibody (Cat#AM8442b). AM8442b was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Western blot analysis of lysates from HeLa, MCF-7, mouse C2C12, mouse NIH/3T3, rat C6 cell line (from left to right), using RPTOR Antibody (Cat. #AM8442b). AM8442b was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L (HRP) at 1:10000 dilution was used as the secondary antibody. Lysates at 20µg per lane.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.