

# CHRM2 Antibody

Purified Mouse Monoclonal Antibody (Mab)  
Catalog # AM8445b

## Product Information

---

<b>Application</b>	WB, IF, IHC-P, FC
<b>Primary Accession</b>	<a href="#">P08172</a>
<b>Reactivity</b>	Human, Mouse
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1, $\kappa$
<b>Clone Names</b>	1424CT461.78.60
<b>Calculated MW</b>	51715

## Additional Information

---

<b>Gene ID</b>	1129
<b>Other Names</b>	Muscarinic acetylcholine receptor M2, CHRM2
<b>Target/Specificity</b>	This antibody is generated from a mouse immunized with a recombinant protein.
<b>Dilution</b>	FC~~1:25 IF~~1:25 IHC-P~~1:100~500 WB~~1:500
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	CHRM2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	CHRM2
<b>Function</b>	The muscarinic acetylcholine receptor mediates various cellular responses, including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is adenylate cyclase inhibition. Signaling promotes

phospholipase C activity, leading to the release of inositol trisphosphate (IP3); this then triggers calcium ion release into the cytosol.

## Cellular Location

Cell membrane; Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane; Multi-pass membrane protein  
Note=Phosphorylation in response to agonist binding promotes receptor internalization. {ECO:0000250 | UniProtKB:P06199}


## Background

The muscarinic acetylcholine receptor mediates various cellular responses, including inhibition of adenylate cyclase, breakdown of phosphoinositides and modulation of potassium channels through the action of G proteins. Primary transducing effect is adenylate cyclase inhibition.


## References

Bonner T.I., et al. *Science* 237:527-532(1987).  
Peralta E.G., et al. *EMBO J.* 6:3923-3929(1987).  
Puhl H.L. III, et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.  
Kitano T., et al. *Mol. Biol. Evol.* 21:936-944(2004).  
Gurevich V.V., et al. *J. Biol. Chem.* 270:720-731(1995).

## Images





Overlay histogram showing SH-SY5Y cells stained with (green line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (1:25 dilution) for 60 min at 37°C. The secondary antibody used was Alexa Fluor® 488 goat anti-mouse IgG (166821) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG1 (1 µg/1x10<sup>6</sup> cells) used under the same conditions. Acquisition of >10,000 events was performed.




Fluorescent image of SH-SY5Y cells stained with CHRM2 Antibody (Cat#AM8445b). AM8445b was diluted at 1:25 dilution. An Alexa Fluor® 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue).

Immunohistochemical analysis of paraffin-embedded H. brain section using CHRM2 (Cat#AM8445b). AM8445b was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Immunohistochemical analysis of paraffin-embedded H. heart section using CHR2 (Cat#AM8445b ). AM8445b was diluted at 1:25 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.



Western blot analysis of lysates from SH-SY5Y cell line, human brain, mouse brain tissue(from left to right), using CHR2 Antibody(Cat. #AM8445b). AM8445b was diluted at 1:500 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 20µg per lane.