

# HINT1 Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AM8470b

#### **Product Information**

Application	WB, FC, IHC-P, IF, E
Primary Accession	<u>P49773</u>
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1500CT836.13.93
Calculated MW	13802

### **Additional Information**

Gene ID	3094
Other Names	Histidine triad nucleotide-binding protein 1, 3, Adenosine 5'-monophosphoramidase, Protein kinase C inhibitor 1, Protein kinase C-interacting protein 1, PKCI-1, HINT1, HINT, PKCI1, PRKCNH1
Target/Specificity	This HINT1 antibody is generated from a mouse immunized with a recombinant protein of human HINT1.
Dilution	WB~~1:1000 FC~~1:25 IHC-P~~1:100~500 IF~~1:25 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HINT1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### **Protein Information**

Name	HINT1
Synonyms	HINT, PKCI1, PRKCNH1
Function	Exhibits adenosine 5'-monophosphoramidase activity, hydrolyzing purine nucleotide phosphoramidates with a single phosphate group such as

	adenosine 5'monophosphoramidate (AMP-NH2) to yield AMP and NH2
	(PubMed: <u>15703176</u> , PubMed: <u>16835243</u> , PubMed: <u>17217311</u> ,
	PubMed: <u>17337452</u> , PubMed: <u>22329685</u> , PubMed: <u>23614568</u> ,
	PubMed: <u>28691797</u> , PubMed: <u>29787766</u> , PubMed: <u>31990367</u> ). Hydrolyzes
	adenosine 5'monophosphomorpholidate (AMP-morpholidate) and guanosine
	5'monophosphomorpholidate (GMP-morpholidate) (PubMed: <u>15703176</u> ,
	PubMed: <u>16835243</u> ). Hydrolyzes lysyl-AMP (AMP-N-epsilon-(N-alpha-acetyl
	lysine methyl ester)) generated by lysine tRNA ligase, as well as Met- AMP,
	His-AMP and Asp-AMP, lysyl-GMP (GMP-N-epsilon-(N-alpha-acetyl lysine
	methyl ester)) and AMP-N-alanine methyl ester (PubMed: <u>15703176</u> ,
	PubMed: <u>17337452</u> , PubMed: <u>22329685</u> ). Hydrolyzes 3-indolepropionic acyl-
	adenylate, tryptamine adenosine phosphoramidate monoester and other
	fluorogenic purine nucleoside tryptamine phosphoramidates in vitro
	(PubMed: <u>17217311</u> , PubMed: <u>17337452</u> , PubMed: <u>23614568</u> ,
	PubMed: <u>28691797</u> , PubMed: <u>29787766</u> , PubMed: <u>31990367</u> ). Can also convert
	adenosine 5'-O- phosphorothioate and guanosine 5'-O-phosphorothioate to
	the corresponding nucleoside 5'-O-phosphates with concomitant release of
	hydrogen sulfide (PubMed: <u>30772266</u> ). In addition, functions as scaffolding
	protein that modulates transcriptional activation by the LEF1/TCF1-CTNNB1
	complex and by the complex formed with MITF and CTNNB1
	(PubMed: <u>16014379</u> , PubMed: <u>22647378</u> ). Modulates p53/TP53 levels and
	p53/TP53-mediated apoptosis (PubMed: <u>16835243</u> ). Modulates proteasomal
	degradation of target proteins by the SCF (SKP2-CUL1-F-box protein) E3
	ubiquitin-protein ligase complex (PubMed: <u>19112177</u> ). Also exhibits SUMO-
	specific isopeptidase activity, deconjugating SUMO1 from RGS17
	(PubMed: <u>31088288</u> ). Deconjugates SUMO1 from RANGAP1 (By similarity).
Cellular Location	Cytoplasm. Nucleus. Note=Interaction with CDK7 leads to a more nuclear
	localization.
Tissue Location	Widely expressed

## Background

Hydrolyzes purine nucleotide phosphoramidates with a single phosphate group, including adenosine 5'monophosphoramidate (AMP-NH2), adenosine 5'monophosphomorpholidate (AMP-morpholidate) and guanosine 5'monophosphomorpholidate (GMP-morpholidate). Hydrolyzes lysyl-AMP (AMP-N-epsilon-(N-alpha-acetyl lysine methyl ester)) generated by lysine tRNA ligase, as well as Met-AMP, His- AMP and Asp-AMP, lysyl-GMP (GMP-N-epsilon-(N-alpha-acetyl lysine methyl ester)) and AMP-N-alanine methyl ester. Can also convert adenosine 5'-O-phosphorothioate and guanosine 5'-O- phosphorothioate to the corresponding nucleoside 5'-O-phosphates with concomitant release of hydrogen sulfide. In addition, functions as scaffolding protein that modulates transcriptional activation by the LEF1/TCF1-CTNNB1 complex and by the complex formed with MITF and CTNNB1. Modulates p53/TP53 levels and p53/TP53-mediated apoptosis. Modulates proteasomal degradation of target proteins by the SCF (SKP2-CUL1-F-box protein) E3 ubiquitin- protein ligase complex.

## References

Brzoska P.M.,et al.Genomics 36:151-156(1996). Brzoska P.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 92:7824-7828(1995). Ota T.,et al.Nat. Genet. 36:40-45(2004). Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Lima C.D.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:5357-5362(1996).

#### Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.