

RASSF2 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8525b

Product Information

Application	WB, FC, E
Primary Accession	P50749
Reactivity	Human, Rat, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG2a,k
Clone Names	1509CT269.10.9.29
Calculated MW	37790

Additional Information

Gene ID	9770
Other Names	Ras association domain-containing protein 2, RASSF2, KIAA0168
Target/Specificity	This RASSF2 antibody is generated from a mouse immunized with a recombinant protein between 1-326 amino acids from human RASSF2.
Dilution	WB~~1:2000 FC~~1:25 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RASSF2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RASSF2
Synonyms	CENP-34 {ECO:0000303 PubMed:20813266}, K
Function	Potential tumor suppressor. Acts as a KRAS-specific effector protein. May promote apoptosis and cell cycle arrest. Stabilizes STK3/MST2 by protecting it from proteasomal degradation.
Cellular Location	Nucleus. Cytoplasm. Chromosome, centromere, kinetochore.

Note=Translocates to the cytoplasm in the presence of STK3/MST2 and STK4/MST1

Tissue Location

Widely expressed with highest levels in brain, placenta, peripheral blood and lung. Frequently down-regulated in lung tumor cell lines.


Background

Potential tumor suppressor. Acts as a KRAS-specific effector protein. May promote apoptosis and cell cycle arrest. Stabilizes STK3/MST2 by protecting it from proteasomal degradation.


References

Burbee D.G.,et al.Submitted (SEP-2002) to the EMBL/GenBank/DDBJ databases.
Nagase T.,et al.DNA Res. 3:17-24(1996).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Suzuki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Images



Overlay histogram showing Raji cells stained with AM8525b(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8525b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OJ192088) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG2a (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.



Anti-RASSF2 Antibody at 1:2000 dilution + Raji whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.