

# PLIN3 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8538b

## Product Information

---

|                   |                        |
|-------------------|------------------------|
| Application       | WB, IF, E              |
| Primary Accession | <a href="#">O60664</a> |
| Reactivity        | Human                  |
| Host              | Mouse                  |
| Clonality         | monoclonal             |
| Isotype           | IgG2a,k                |
| Clone Names       | 1651CT157.85.94        |
| Calculated MW     | 47075                  |

## Additional Information

---

|                    |  |
|--------------------|--|
| Gene ID            | 10226  |
| Other Names        | Perilipin-3, 47 kDa mannose 6-phosphate receptor-binding protein, 47 kDa MPR-binding protein, Cargo selection protein TIP47, Mannose-6-phosphate receptor-binding protein 1, Placental protein 17, PP17, PLIN3, M6PRBP1, TIP47 |
| Target/Specificity | This PLIN3 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 1-434 amino acids from human PLIN3.  |
| Dilution           | WB~~1:2000 IF~~1:25 E~~Use at an assay dependent concentration.  |
| Format             | Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.  |
| Storage            | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.  |
| Precautions        | PLIN3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.   |

## Protein Information

---

|          |   |
|----------|---|
| Name     | PLIN3   |
| Synonyms | M6PRBP1, TIP47 {ECO:0000303   PubMed:95901  |
| Function | Structural component of lipid droplets, which is required for the formation and maintenance of lipid storage droplets (PubMed: <a href="#">34077757</a> ). Required for |

the transport of mannose 6-phosphate receptors (MPR) from endosomes to the trans-Golgi network (PubMed:[9590177](#)).

## Cellular Location

Lipid droplet. Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasm. Note=Membrane associated on endosomes (PubMed:15545278). Detected in the envelope and the core of lipid bodies and in lipid sails (PubMed:15545278)


## Background

Required for the transport of mannose 6-phosphate receptors (MPR) from endosomes to the trans-Golgi network.


## References

Diaz E.,et al.Cell 93:433-443(1998).  
Than N.G.,et al.Eur. J. Biochem. 258:752-757(1998).  
Than N.G.,et al.Tumor Biol. 20:184-192(1999).  
Kalnina N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.  
Ota T.,et al.Nat. Genet. 36:40-45(2004).

## Images



Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized U-2 OS (human osteosarcoma cell line) cells labeling PLIN3 with AM8538b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG (NA166821) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on U-2 OS cell line. The nuclear counter stain is DAPI (blue).



All lanes : Anti-PLIN3 Antibody at 1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: K562 whole cell lysate Lane 3: Daudi whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.