

RAB20 Antibody [Knockdown Validated]

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8559b

Product Information

Application	WB, E
Primary Accession	Q9NX57
Reactivity	Human, Mouse
Predicted	Canine
Host	Mouse
Clonality	monoclonal
Isotype	IgG2a
Clone Names	1694CT210.142.16
Calculated MW	26277

Additional Information

Gene ID	55647
Other Names	Ras-related protein Rab-20, RAB20
Target/Specificity	This RAB20 antibody is generated from a mouse immunized with recombinant protein from human RAB20.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RAB20 Antibody [Knockdown Validated] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RAB20 (HGNC:18260)
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (By similarity). RAB20 plays a role in apical

endocytosis/recycling. Plays a role in the maturation and acidification of phagosomes that engulf pathogens, such as *S.aureus* and *M.tuberculosis*. Plays a role in the fusion of phagosomes with lysosomes.

Cellular Location

Golgi apparatus. Cytoplasmic vesicle, phagosome Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Note=Highly enriched on apical endocytic structures in polarized epithelial cells of kidney proximal tubules (By similarity). Recruited to phagosomes containing *S.aureus* or *M.tuberculosis* (PubMed:21255211) {ECO:0000250|UniProtKB:P35295, ECO:0000269|PubMed:21255211}

Tissue Location

Low or absent expression in normal pancreas and stronger expression in 15 of 18 exocrine pancreatic adenocarcinomas (at protein level).


Background

Plays a role in apical endocytosis/recycling. Plays a role in the maturation and acidification of phagosomes that engulf pathogens, such as *S.aureus* and *M.tuberculosis*. Plays a role in the fusion of phagosomes with lysosomes.


References

Amillet J.-M.,et al.Hum. Pathol. 37:256-263(2006).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Dunham A.,et al.Nature 428:522-528(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Seto S.,et al.Traffic 12:407-420(2011).

Images



All lanes : Anti-RAB20 Antibody at 1:2000-1:4000 dilution
Lane 1: HeLa whole cell lysate Lane 2: MCF-7 whole cell lysate
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-RAB20 Antibody at 1:2000 dilution
Lane 1: HepG2 Lane 2: HepG2-Knockdown
Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.