

CTCF Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8583b

Product Information

Application	WB, E
Primary Accession	P49711
Reactivity	Human, Rat, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1723CT382.14.9
Calculated MW	82785

Additional Information

Gene ID	10664
Other Names	Transcriptional repressor CTCF, 11-zinc finger protein, CCCTC-binding factor, CTCFL paralog, CTCF
Target/Specificity	This CTCF antibody is generated from a mouse immunized with a recombinant protein between 445-727 amino acids from human CTCF.
Dilution	WB~~1:4000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	CTCF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	CTCF
Function	Chromatin binding factor that binds to DNA sequence specific sites and regulates the 3D structure of chromatin (PubMed: 18347100 , PubMed: 18654629 , PubMed: 19322193). Binds together strands of DNA, thus forming chromatin loops, and anchors DNA to cellular structures, such as the nuclear lamina (PubMed: 18347100 , PubMed: 18654629 , PubMed: 19322193). Defines the boundaries between active and heterochromatic DNA via binding

to chromatin insulators, thereby preventing interaction between promoter and nearby enhancers and silencers (PubMed:[18347100](#), PubMed:[18654629](#), PubMed:[19322193](#)). Plays a critical role in the epigenetic regulation (PubMed:[16949368](#)). Participates in the allele-specific gene expression at the imprinted IGF2/H19 gene locus (PubMed:[16107875](#), PubMed:[16815976](#), PubMed:[17827499](#)). On the maternal allele, binding within the H19 imprinting control region (ICR) mediates maternally inherited higher- order chromatin conformation to restrict enhancer access to IGF2 (By similarity). Mediates interchromosomal association between IGF2/H19 and WSB1/NF1 and may direct distant DNA segments to a common transcription factory (By similarity). Regulates asynchronous replication of IGF2/H19 (By similarity). Plays a critical role in gene silencing over considerable distances in the genome (By similarity). Preferentially interacts with unmethylated DNA, preventing spreading of CpG methylation and maintaining methylation-free zones (PubMed:[18413740](#)). Inversely, binding to target sites is prevented by CpG methylation (PubMed:[18413740](#)). Plays an important role in chromatin remodeling (PubMed:[18413740](#)). Can dimerize when it is bound to different DNA sequences, mediating long-range chromatin looping (PubMed:[12191639](#)). Causes local loss of histone acetylation and gain of histone methylation in the beta-globin locus, without affecting transcription (PubMed:[12191639](#)). When bound to chromatin, it provides an anchor point for nucleosomes positioning (PubMed:[12191639](#)). Seems to be essential for homologous X-chromosome pairing (By similarity). May participate with Tsix in establishing a regulatable epigenetic switch for X chromosome inactivation (PubMed:[11743158](#)). May play a role in preventing the propagation of stable methylation at the escape genes from X-inactivation (PubMed:[11743158](#)). Involved in sister chromatid cohesion (PubMed:[12191639](#)). Associates with both centromeres and chromosomal arms during metaphase and required for cohesin localization to CTCF sites (PubMed:[18550811](#)). Plays a role in the recruitment of CENPE to the pericentromeric/centromeric regions of the chromosome during mitosis (PubMed:[26321640](#)). Acts as a transcriptional repressor binding to promoters of vertebrate MYC gene and BAG1 gene (PubMed:[18413740](#), PubMed:[8649389](#), PubMed:[9591631](#)). Also binds to the PLK and PIM1 promoters (PubMed:[12191639](#)). Acts as a transcriptional activator of APP (PubMed:[9407128](#)). Regulates APOA1/C3/A4/A5 gene cluster and controls MHC class II gene expression (PubMed:[18347100](#), PubMed:[19322193](#)). Plays an essential role in oocyte and preimplantation embryo development by activating or repressing transcription (By similarity). Seems to act as tumor suppressor (PubMed:[12191639](#)).

Cellular Location

Nucleus, nucleoplasm. Chromosome. Chromosome, centromere. Note=May translocate to the nucleolus upon cell differentiation. Associates with both centromeres and chromosomal arms during metaphase. Associates with the H19 ICR in mitotic chromosomes. May be preferentially excluded from heterochromatin during interphase

Tissue Location

Ubiquitous. Absent in primary spermatocytes.

Background


Chromatin binding factor that binds to DNA sequence specific sites. Involved in transcriptional regulation by binding to chromatin insulators and preventing interaction between promoter and nearby enhancers and silencers. Acts as transcriptional repressor binding to promoters of vertebrate MYC gene and BAG1 gene. Also binds to the PLK and PIM1 promoters. Acts as a transcriptional activator of APP. Regulates APOA1/C3/A4/A5 gene cluster and controls MHC class II gene expression. Plays an essential role in oocyte and preimplantation embryo development by activating or repressing transcription. Seems to act as tumor suppressor. Plays a critical role in the epigenetic regulation. Participates in the allele-specific gene expression at the imprinted IGF2/H19 gene locus. On the maternal allele, binding within the H19 imprinting control region (ICR) mediates maternally inherited higher-order chromatin conformation to restrict

enhancer access to IGF2. Plays a critical role in gene silencing over considerable distances in the genome. Preferentially interacts with unmethylated DNA, preventing spreading of CpG methylation and maintaining methylation-free zones. Inversely, binding to target sites is prevented by CpG methylation. Plays a important role in chromatin remodeling. Can dimerize when it is bound to different DNA sequences, mediating long-range chromatin looping. Mediates interchromosomal association between IGF2/H19 and WSB1/NF1 and may direct distant DNA segments to a common transcription factory. Causes local loss of histone acetylation and gain of histone methylation in the beta-globin locus, without affecting transcription. When bound to chromatin, it provides an anchor point for nucleosomes positioning. Seems to be essential for homologous X-chromosome pairing. May participate with Tsix in establishing a regulatable epigenetic switch for X chromosome inactivation. May play a role in preventing the propagation of stable methylation at the escape genes from X- inactivation. Involved in sister chromatid cohesion. Associates with both centromeres and chromosomal arms during metaphase and required for cohesin localization to CTCF sites. Regulates asynchronous replication of IGF2/H19.

References

Filippova G.N.,et al.Mol. Cell. Biol. 16:2802-2813(1996).
 Filippova G.N.,et al.Genes Chromosomes Cancer 22:26-36(1998).
 Filippova G.N.,et al.Cancer Res. 62:48-52(2002).
 Kalnine N.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.
 Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-CTCF Antibody at 1:4000 dilution Lane 1: K562 whole cell lysate Lane 2: Jurkat whole cell lysate Lane 3: HeLa whole cell lysate Lane 4: C6 whole cell lysate Lane 5: MOLT-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 83 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.