

PACSIN2 Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AM8589b

Product Information

Application WB, E **Primary Accession** Q9UNFO

Reactivity Human, Mouse

HostMouseClonalitymonoclonalIsotypeIgG1,k

Clone Names 1735CT116.76.60

Calculated MW 55739

Additional Information

Gene ID 11252

Other Names Protein kinase C and casein kinase substrate in neurons protein 2,

Syndapin-2, Syndapin-II, PACSIN2

Target/Specificity This PACSIN2 antibody is generated from a mouse immunized with a

recombinant protein between 250-486 amino acids from human PACSIN2.

Dilution WB~~1:2000 E~~Use at an assay dependent concentration.

Format Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide.

This antibody is purified through a protein G column, followed by dialysis

against PBS.

Storage Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store

at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions PACSIN2 Antibody is for research use only and not for use in diagnostic or

therapeutic procedures.

Protein Information

Name PACSIN2

Function Regulates the morphogenesis and endocytosis of caveolae (By similarity).

Lipid-binding protein that is able to promote the tubulation of the

phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus (PubMed:21693584, PubMed:23129763,

PubMed:<u>23236520</u>, PubMed:<u>23596323</u>). Essential for endothelial organization in sprouting angiogenesis, modulates CDH5-based junctions. Facilitates endothelial front-rear polarity during migration by recruiting EHD4 and MICALL1 to asymmetric adherens junctions between leader and follower cells (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9WVE8}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q9WVE8}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q9WVE8}; Peripheral membrane protein {ECO:0000250|UniProtKB:O9WVE8}: Cvtoplasmic side {ECO:0000250|UniProtKB:Q9WVE8}. Cell projection, ruffle membrane {ECO:0000250|UniProtKB:Q9WVE8}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9WVE8}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9WVE8}. Early endosome {ECO:0000250|UniProtKB:Q9WVE8}. Recycling endosome membrane. Cell membrane {ECO:0000250|UniProtKB:Q9WVE8}; Peripheral membrane protein {ECO:0000250 | UniProtKB:Q9WVE8}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9WVE8}. Cell projection. Membrane, caveola. Cell junction, adherens junction {ECO:0000250 | UniProtKB:Q9WVE8}. Note=Detected at the neck of flask- shaped caveolae. Localization to tubular recycling endosomes probably requires interaction with MICALL1 and EHD1 {ECO:0000250|UniProtKB:Q9WVE8}

Tissue Location

Widely expressed.


Background

Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle- mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus. May also play a role in the formation of caveolae at the cell membrane. Recruits DNM2 to caveolae, and thereby plays a role in caveola-mediated endocytosis.

References

Ritter B., et al. FEBS Lett. 454:356-362(1999). Wiemann S., et al. Genome Res. 11:422-435(2001). Collins J.E., et al. Genome Biol. 5:R84.1-R84.11(2004). Dunham I., et al. Nature 402:489-495(1999). Modregger J., et al. J. Cell Sci. 113:4511-4521(2000).

Images



All lanes: Anti-PACSIN2 Antibody at 1:2000 dilution Lane 1: Daudi whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: human brain lysate Lane 4: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size: 56 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.