

FNDC5 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8591b

Product Information

Application	WB, E
Primary Accession	Q8NAU1
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1736CT532.85.53.92
Calculated MW	28283

Additional Information

Gene ID	252995
Other Names	Fibronectin type III domain-containing protein 5, Fibronectin type III repeat-containing protein 2, Irisin, FNDC5, FRCP2
Target/Specificity	This FNDC5 antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 12-152 amino acids from human FNDC5.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	FNDC5 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	FNDC5
Synonyms	FRCP2
Function	[Irisin]: Mediates beneficial effects of muscular exercise. Induces browning of white adipose tissue by stimulating UCP1 expression, at least in part, via the nuclear receptor PPARA.

Cellular Location	[Fibronectin type III domain-containing protein 5]: Cell membrane; Single-pass membrane protein. Peroxisome membrane {ECO:0000250 UniProtKB:Q8K4Z2}; Single-pass membrane protein. Note=Imported into peroxisomes through the PEX5 receptor pathway. {ECO:0000250 UniProtKB:Q8K4Z2}
Tissue Location	Widely expressed, with highest levels in heart. Very low expression, if any, in colon, pancreas and spleen


Background

Irisin: Contrary to mouse, may not be involved in the beneficial effects of muscular exercise, nor in the induction of browning of human white adipose tissue.

References

Ota T.,et al.Nat. Genet. 36:40-45(2004).
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Bostrom P.,et al.Nature 481:463-468(2012).

Images



All lanes: Anti-FND5 Antibody at 1:1000 dilution Lane 1: Mouse skeletal muscle lysate Lane 2: Rat skeletal muscle lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Observed band size: 24 KDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.