

# MMP1 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8598b

## Product Information

---

Application	WB, E
Primary Accession	<a href="#">P03956</a>
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG2a
Clone Names	1054CT20.4.3
Calculated MW	54007

## Additional Information

---

Gene ID	4312
Other Names	Interstitial collagenase, 3.4.24.7, Fibroblast collagenase, Matrix metalloproteinase-1, MMP-1, 22 kDa interstitial collagenase, 27 kDa interstitial collagenase, MMP1, CLG
Target/Specificity	This antibody is generated from a mouse immunized with a recombinant protein from human.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MMP1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	MMP1
Synonyms	CLG
Function	Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X (PubMed: <a href="#">1645757</a> , PubMed: <a href="#">2153297</a> , PubMed: <a href="#">2557822</a> ). In case of HIV infection, interacts and cleaves the secreted

viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity (PubMed:[16807369](#)).

## Cellular Location

Secreted, extracellular space, extracellular matrix


## Background

Cleaves collagens of types I, II, and III at one site in the helical domain. Also cleaves collagens of types VII and X. In case of HIV infection, interacts and cleaves the secreted viral Tat protein, leading to a decrease in neuronal Tat's mediated neurotoxicity.

## References

Templeton N.S.,et al.Cancer Res. 50:5431-5437(1990).  
Whitham S.E.,et al.Biochem. J. 240:913-916(1986).  
Goldberg G.I.,et al.J. Biol. Chem. 261:6600-6605(1986).  
Lin D.,et al.Submitted (DEC-1996) to the EMBL/GenBank/DDBJ databases.  
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

## Images



All lanes : Anti-MMP1 Antibody at 1:2000 dilution Lane 1: Raji whole cell lysate Lane 2: PC-3 whole cell lysate Lane 3: MDA-MB-231 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52/55 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.