

HK2 (Hexokinase II) Antibody [Knockdown Validated]

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8606b

Product Information

Application	WB, IHC-P, E
Primary Accession	P52789
Reactivity	Human
Predicted	Mouse, Rat, Bovine, Canine, Rabbit, Chicken
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1798CT748.15.25
Calculated MW	102380

Additional Information

Gene ID	3099
Other Names	Hexokinase-2, 2.7.1.1, Hexokinase type II, HK II, Muscle form hexokinase, HK2
Target/Specificity	This HK2 (Hexokinase II) antibody is generated from a mouse immunized with a recombinant protein between 1-170 amino acids from human HK2 (Hexokinase II).
Dilution	WB~~1:2000 IHC-P~~1:400 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	HK2 (Hexokinase II) Antibody [Knockdown Validated] is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	HK2 (HGNC:4923)
Function	Catalyzes the phosphorylation of hexose, such as D-glucose and D-fructose, to hexose 6-phosphate (D-glucose 6-phosphate and D- fructose 6-phosphate, respectively) (PubMed: 23185017 , PubMed: 26985301 , PubMed: 29298880). Mediates the initial step of glycolysis by catalyzing phosphorylation of D-glucose to D-glucose 6-phosphate (PubMed: 29298880). Plays a key role in

maintaining the integrity of the outer mitochondrial membrane by preventing the release of apoptogenic molecules from the intermembrane space and subsequent apoptosis (PubMed:[18350175](#)).

Cellular Location

Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasm, cytosol Note=The mitochondrial-binding peptide (MBP) region promotes association with the mitochondrial outer membrane (PubMed:29298880) The interaction with the mitochondrial outer membrane via the mitochondrial-binding peptide (MBP) region promotes higher stability of the protein (PubMed:29298880). Release from the mitochondrial outer membrane into the cytosol induces permeability transition pore (PTP) opening and apoptosis (PubMed:18350175).


Tissue Location

Predominant hexokinase isozyme expressed in insulin-responsive tissues such as skeletal muscle


References

Deeb S.S.,et al.Biochem. Biophys. Res. Commun. 197:68-74(1993).
Lehto M.,et al.Diabetologia 38:1466-1474(1995).
Malkki M.,et al.Submitted (MAY-1999) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Shinohara Y.,et al.Cancer Lett. 82:27-32(1994).

Images




All lanes : Anti-HK2 (Hexokinase II) Antibody at 1:500-1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: 293 whole cell lysate Lane 3: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 102 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Immunohistochemical analysis of paraffin-embedded Human Skeletal muscle section using Pink1(Cat#AM8606b). AM8606b was diluted at 1:400 dilution. A undiluted biotinylated goat polyvalent antibody was used as the secondary, followed by DAB staining.

All lanes : Anti-HK2 (Hexokinase II) Antibody at 1:2000 dilution Lane 1: HeLa Lane 2: HeLa-Knockdown Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613)

at 1/8000 dilution.



Citations

- [Dickkopf-related protein 3 alters aerobic glycolysis in pancreatic cancer BxPC-3 cells, promoting CD4 T-cell activation and function](#)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.