

ADK Antibody (N-term)

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8619b

Product Information

Application	WB, IHC-P, E
Primary Accession	P55263
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1103CT3.4.3
Calculated MW	40545

Additional Information

Gene ID	132
Other Names	Adenosine kinase, AK, 2.7.1.20, Adenosine 5'-phosphotransferase, ADK
Target/Specificity	This ADK antibody is generated from a mouse immunized with a recombinant protein between 1-345 amino acids from human ADK.
Dilution	WB~~1:4000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	ADK Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	ADK (HGNC:257)
Function	Catalyzes the phosphorylation of the purine nucleoside adenosine at the 5' position in an ATP-dependent manner. Serves as a potential regulator of concentrations of extracellular adenosine and intracellular adenine nucleotides.
Cellular Location	[Isoform 1]: Nucleus

Tissue Location

Widely expressed. Highest level in placenta, liver, muscle and kidney.

Background

ATP dependent phosphorylation of adenosine and other related nucleoside analogs to monophosphate derivatives. Serves as a potential regulator of concentrations of extracellular adenosine and intracellular adenine nucleotides.

References

Spychala J.,et al.Proc. Natl. Acad. Sci. U.S.A. 93:1232-1237(1996).


Singh B.,et al.Eur. J. Biochem. 241:564-571(1996).

McNally T.,et al.Biochem. Biophys. Res. Commun. 231:645-650(1997).

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Deloukas P.,et al.Nature 429:375-381(2004).

Images



All lanes: Anti-ADK Antibody (N-term) at 1:2000 dilution
Lane 1: HL-60 whole cell lysate Lane 2: A431 whole cell lysate Lane 3: 293T whole cell lysate Lane 4: MCF-7 whole cell lysate Lane 5: Mouse liver lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Mouse IgG, (H+L), Peroxidase conjugated (ASP1613) at 1/8000 dilution. Observed band size: 45 KDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.