

# MBD2 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8622b

## Product Information

---

<b>Application</b>	WB, IHC-P, E
<b>Primary Accession</b>	<a href="#">Q9UBB5</a>
<b>Reactivity</b>	Human, Rat, Mouse
<b>Host</b>	Mouse
<b>Clonality</b>	monoclonal
<b>Isotype</b>	IgG1,k
<b>Clone Names</b>	1812CT625.9.45
<b>Calculated MW</b>	43255

## Additional Information

---

<b>Gene ID</b>	8932
<b>Other Names</b>	Methyl-CpG-binding domain protein 2, Demethylase, DMTase, Methyl-CpG-binding protein MBD2, MBD2
<b>Target/Specificity</b>	This MBD2 antibody is generated from a mouse immunized with a recombinant protein between 10-228 amino acids from human MBD2.
<b>Dilution</b>	WB~~1:2000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration.
<b>Format</b>	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
<b>Storage</b>	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
<b>Precautions</b>	MBD2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

<b>Name</b>	MBD2 ( <a href="#">HGNC:6917</a> )
<b>Function</b>	Binds CpG islands in promoters where the DNA is methylated at position 5 of cytosine within CpG dinucleotides (PubMed: <a href="#">9774669</a> ). Binds hemimethylated DNA as well (PubMed: <a href="#">10947852</a> , PubMed: <a href="#">24307175</a> ). Recruits histone deacetylases and DNA methyltransferases to chromatin (PubMed: <a href="#">10471499</a> , PubMed: <a href="#">10947852</a> ). Acts as a component of the histone deacetylase NuRD complex which participates in the remodeling of chromatin

(PubMed:[16428440](#), PubMed:[28977666](#)). Acts as a transcriptional repressor and plays a role in gene silencing (PubMed:[10471499](#), PubMed:[10947852](#), PubMed:[16415179](#)). Functions as a scaffold protein, targeting GATAD2A and GATAD2B to chromatin to promote repression (PubMed:[16415179](#)). May enhance the activation of some unmethylated cAMP-responsive promoters (PubMed:[12665568](#)).

#### Cellular Location

Nucleus. Chromosome Note=Nuclear, in discrete foci (PubMed:12183469). Detected at replication foci in late S phase. Localizes to methylated chromatin (PubMed:16428440). Localizes to sites of DNA damage in a manner partially dependent on ZMYND8 (PubMed:27732854)

#### Tissue Location

Highly expressed in brain, heart, kidney, stomach, testis and placenta.

## Background

---

Binds CpG islands in promoters where the DNA is methylated at position 5 of cytosine within CpG dinucleotides. Binds hemimethylated DNA as well. Recruits histone deacetylases and DNA methyltransferases. Acts as transcriptional repressor and plays a role in gene silencing. Functions as a scaffold protein, targeting GATAD2A and GATAD2B to chromatin to promote repression. May enhance the activation of some unmethylated cAMP-responsive promoters.


## References

---

Hendrich B.,et al.Mol. Cell. Biol. 18:6538-6547(1998).  
Hendrich B.,et al.Mamm. Genome 10:906-912(1999).  
Bhattacharya S.K.,et al.Nature 397:579-583(1999).  
Ng H.-H.,et al.Nat. Genet. 23:58-61(1999).  
Tatematsu K.,et al.Genes Cells 5:677-688(2000).


## Images

---



AM8622b staining MBD2 in human colon tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

All lanes : Anti-MBD2 Antibody at 1:2000 dilution Lane 1: K562 whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 43 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.