

DKK2 Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AM8643b

Product Information

Application	WB, E
Primary Accession	<u>Q9UBU2</u>
Reactivity	Human, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1833CT856.7.20
Calculated MW	28447

Additional Information

Gene ID	27123
Other Names	Dickkopf-related protein 2, Dickkopf-2, Dkk-2, hDkk-2, DKK2
Target/Specificity	This DKK2 antibody is generated from a mouse immunized with a recombinant protein of human DKK2.
Dilution	WB~~1:2000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	DKK2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

NameDKK2FunctionAntagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with
Wnt and by forming a ternary complex with the transmembrane protein
KREMEN that promotes internalization of LRP5/6. DKKs play an important role
in vertebrate development, where they locally inhibit Wnt regulated processes
such as antero-posterior axial patterning, limb development, somitogenesis
and eye formation. In the adult, Dkks are implicated in bone formation and
bone disease, cancer and Alzheimer disease (By similarity).

Cellular Location	Secreted.
Tissue Location	Expressed in heart, brain, skeletal muscle and lung

Background

Antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero- posterior axial patterning, limb development, somitogenesis and eye formation. In the adult, Dkks are implicated in bone formation and bone disease, cancer and Alzheimer disease (By similarity).

References

Krupnik V.E.,et al.Gene 238:301-313(1999). Tanaka S.,et al.Submitted (OCT-1999) to the EMBL/GenBank/DDBJ databases. Clark H.F.,et al.Genome Res. 13:2265-2270(2003). Ota T.,et al.Nat. Genet. 36:40-45(2004). Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



All lanes : Anti-DKK2 Antibody at 1:2000 dilution Lane 1: Human brain lysate Lane 2: Mouse heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.