

RAB13 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8693b

Product Information

Application	WB, E
Primary Accession	P51153
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1
Clone Names	1600CT724.2.34
Calculated MW	22774

Additional Information

Gene ID	5872
Other Names	Ras-related protein Rab-13, Cell growth-inhibiting gene 4 protein, RAB13
Target/Specificity	This RAB13 antibody is generated from a mouse immunized with a recombinant protein between 1-203 amino acids from human RAB13.
Dilution	WB~~1:1000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	RAB13 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	RAB13 (HGNC:9762)
Function	The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB13 is involved in endocytic recycling and regulates the transport to the plasma membrane of transmembrane proteins like the

tight junction protein OCLN/occludin. Thereby, it regulates the assembly and the activity of tight junctions. Moreover, it may also regulate tight junction assembly by activating the PKA signaling pathway and by reorganizing the actin cytoskeleton through the activation of the downstream effectors PRKACA and MICALL2 respectively. Through its role in tight junction assembly, may play a role in the establishment of Sertoli cell barrier. Plays also a role in angiogenesis through regulation of endothelial cells chemotaxis. Also involved in neurite outgrowth. Has also been proposed to play a role in post-Golgi membrane trafficking from the TGN to the recycling endosome. Finally, it has been involved in insulin-induced transport to the plasma membrane of the glucose transporter GLUT4 and therefore may play a role in glucose homeostasis.

Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic vesicle membrane; Lipid-anchor; Cytoplasmic side. Cell junction, tight junction. Golgi apparatus, trans-Golgi network membrane Recycling endosome membrane. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q9DD03}. Note=Tight junctions or associated with vesicles scattered throughout the cytoplasm in cells lacking tight junctions (PubMed:8294494) Relocalizes to the leading edge of lamellipodia in migrating endothelial cells (By similarity). {ECO:0000250|UniProtKB:Q9DD03, ECO:0000269|PubMed:8294494}

Tissue Location

Detected in several types of epithelia, including intestine, kidney, liver and in endothelial cells

Background


The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. That Rab is involved in endocytic recycling and regulates the transport to the plasma membrane of transmembrane proteins like the tight junction protein OCLN/occludin. Thereby, it regulates the assembly and the activity of tight junctions. Moreover, it may also regulate tight junction assembly by activating the PKA signaling pathway and by reorganizing the actin cytoskeleton through the activation of the downstream effectors PRKACA and MICALL2 respectively. Through its role in tight junction assembly, may play a role in the establishment of Sertoli cell barrier. Plays also a role in angiogenesis through regulation of endothelial cells chemotaxis. Also involved in neurite outgrowth. Has also been proposed to play a role in post-Golgi membrane trafficking from the TGN to the recycling endosome. Finally, it has been involved in insulin- induced transport to the plasma membrane of the glucose transporter GLUT4 and therefore may play a role in glucose homeostasis.

References

Zahraoui A.,et al.J. Cell Biol. 124:101-115(1994).
Kim J.W.,et al.Submitted (SEP-2003) to the EMBL/GenBank/DDBJ databases.
Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Kalnina N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.

Images

All lanes : Anti-RAB13 Antibody at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: A431 whole cell lysate Lane 3: A549 whole cell lysate Lane 4: U-87 MG whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat



Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.