

GAPDH Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8699b

Product Information

Application	WB, E
Primary Accession	P04406
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Clone Names	1653CT492.29.66
Calculated MW	36053

Additional Information

Gene ID	2597
Other Names	Glyceraldehyde-3-phosphate dehydrogenase, GAPDH, 1.2.1.12, Peptidyl-cysteine S-nitrosylase GAPDH, 2.6.99.-, GAPDH, GAPD
Target/Specificity	This GAPDH antibody is generated from a mouse immunized with a recombinant protein between 43-335 amino acids from human GAPDH.
Dilution	WB~~1:8000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	GAPDH Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	GAPDH {ECO:0000303 PubMed:2987855, ECO:0000312 HGNC:HGNC:4141}
Function	Catalyzes the conversion of D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate in glycolysis and the reverse reaction in gluconeogenesis (PubMed: 11724794 , PubMed: 3170585). Also shows nitrosylase activity, thereby playing a role in nuclear functions (PubMed: 11724794 , PubMed: 3170585). Modulates the organization and assembly of the cytoskeleton (By similarity). Facilitates the CHP1- dependent

microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes (PubMed:[23071094](#)). Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruloplasmin) and suppresses their translation (PubMed:[23071094](#)). Also plays a role in innate immunity by promoting TNF-induced NF-kappa-B activation and type I interferon production, via interaction with TRAF2 and TRAF3, respectively (PubMed:[23332158](#), PubMed:[27387501](#)). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis (By similarity). Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).

Cellular Location

Cytoplasm, cytosol. Nucleus {ECO:0000250 | UniProtKB:P04797}. Cytoplasm, perinuclear region. Membrane Cytoplasm, cytoskeleton {ECO:0000250 | UniProtKB:P04797} Note=Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261) {ECO:0000250 | UniProtKB:P04797, ECO:0000269 | PubMed:12829261}

Background

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC. Modulates the organization and assembly of the cytoskeleton. Facilitates the CHP1-dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate. Component of the GAIT (gamma interferon- activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruloplasmin) and suppresses their translation.


References

- Hanauer A., et al. EMBO J. 3:2627-2633(1984).
Arcari P., et al. Nucleic Acids Res. 12:9179-9189(1984).
Tso J.Y., et al. Nucleic Acids Res. 13:2485-2502(1985).
Tokunaga K., et al. Cancer Res. 47:5616-5619(1987).
Allen R.W., et al. J. Biol. Chem. 262:649-653(1987).

Images

All lanes : Anti-GAPDH Antibody at 1:8000 dilution Lane 1: C6 whole cell lysate Lane 2: mouse brain lysate Lane 3: NIH/3T3 whole cell lysate Lane 4: Hela whole cell lysate Lane 5: Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L),

Peroxidase conjugated at 1/10000 dilution. Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.