

PURB Antibody (C-term)

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AM8708B

Product Information

Application	WB, E
Primary Accession	Q96QR8
Reactivity	Human, Mouse
Predicted	Human, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1, κ
Clone Names	2117CT1176.48.6
Calculated MW	33241

Additional Information

Gene ID	5814
Other Names	Transcriptional activator protein Pur-beta, Purine-rich element-binding protein B, PURB
Target/Specificity	This PURB antibody is generated from a mouse immunized with a recombinant protein of human PURB.
Dilution	WB~~1:8000 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	PURB Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	PURB
Function	Transcriptional regulator which can act as an activator or a repressor. Represses the transcription of ACTA2 in fibroblasts and smooth muscle cells via its ability to interact with the purine-rich strand of a MCAT- containing element in the 5' flanking region of the gene. Represses the transcription of MYOCD, capable of repressing all isoforms of MYOCD but the magnitude of

the repressive effects is most notable for the SMC- specific isoforms. Promotes hepatic glucose production by activating the transcription of ADCY6, leading to cAMP accumulation, increased PKA activity, CREB activation, and increased transcription of PCK1 and G6PC genes (By similarity). Has capacity to bind repeated elements in single-stranded DNA such as the purine-rich single strand of the PUR element located upstream of the MYC gene (PubMed:[1448097](#)). Participates in transcriptional and translational regulation of alpha-MHC expression in cardiac myocytes by binding to the purine-rich negative regulatory (PNR) element Modulates constitutive liver galectin-3 gene transcription by binding to its promoter. May play a role in the dendritic transport of a subset of mRNAs (By similarity).

Cellular Location

Nucleus.

Tissue Location

Expressed in myocardium of heart failure patients.


Background

Has capacity to bind repeated elements in single- stranded DNA such as the purine-rich single strand of the PUR element located upstream of the MYC gene. Plays a role in the control of vascular smooth muscle (VSM) alpha-actin gene transcription as repressor in myoblasts and fibroblasts. Participates in transcriptional and translational regulation of alpha-MHC expression in cardiac myocytes by binding to the purine- rich negative regulatory (PNR) element. Modulates constitutive liver galectin-3 gene transcription by binding to its promoter. May play a role in the dendritic transport of a subset of mRNAs (By similarity).

References

Bergemann A.D.,et al.Mol. Cell. Biol. 12:5673-5682(1992).
 Scherer S.W.,et al.Science 300:767-772(2003).
 Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
 Hillier L.W.,et al.Nature 424:157-164(2003).
 Lezon-Geyda K.,et al.Leukemia 15:954-962(2001).

Images



All lanes : Anti-PURB Antibody (C-term) at 1:8000 dilution
 Lane 1: Jurkat whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: NCI-H460 whole cell lysate Lane 4: HepG2 whole cell lysate Lane 5: MCF-7 whole cell lysate Lane 6: NIH/3T3 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.