

# MAP3K13 (LZK) Antibody (C-term)

Mouse Monoclonal Antibody (Mab)

Catalog # AM9976b

## Product Information

---

Application	WB, IF, E
Primary Accession	<a href="#">O43283</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Clone Names	810CT26.6.1.1
Calculated MW	108296
Antigen Region	52-305

## Additional Information

---

Gene ID	9175
Other Names	Mitogen-activated protein kinase kinase kinase 13, Leucine zipper-bearing kinase, Mixed lineage kinase, MLK, MAP3K13 ( <a href="#">HGNC:6852</a> )
Target/Specificity	Purified His-tagged MAP3K13 protein fragment was used to produced this monoclonal antibody.
Dilution	WB~~1:1000 IF~~1:100 E~~Use at an assay dependent concentration.
Format	Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Euglobin precipitation followed by dialysis against PBS.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MAP3K13 (LZK) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

## Protein Information

---

Name	MAP3K13 ( <a href="#">HGNC:6852</a> )
Function	Activates the JUN N-terminal pathway through activation of the MAP kinase kinase MAP2K7. Acts synergistically with PRDX3 to regulate the activation of NF-kappa-B in the cytosol. This activation is kinase-dependent and involves activating the IKK complex, the IKBKB- containing complex that phosphorylates inhibitors of NF-kappa-B.

**Cellular Location**

Cytoplasm. Membrane; Peripheral membrane protein

**Tissue Location**

Expressed in the adult brain, liver, placenta and pancreas, with expression strongest in the pancreas

## Background

---

Activates the JUN N-terminal pathway through activation of the MAP kinase kinase MAP2K7. Acts synergistically with PRDX3 to regulate the activation of NF-kappa-B in the cytosol. This activation is kinase-dependent and involves activating the IKK complex, the IKBKB-containing complex that phosphorylates inhibitors of NF-kappa-B.


## References

---


Ikeda A., et al. J. Biochem. 130:773-781(2001).  
Masaki M., et al. Eur. J. Biochem. 270:76-83(2003).  
Greenman C., et al. Nature 446:153-158(2007).  
Sakuma H., et al. J. Biol. Chem. 272:28622-28629(1997).  
Ota T., et al. Nat. Genet. 36:40-45(2004).

## Images

---



Anti-MAP3K13 (LZK) Antibody (C-term) at 1:500 dilution + NCI-H460 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgM, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 110 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Fluorescent image of NCI-H460 cells stained with MAP3K13 (LZK) Antibody (N-term)(Cat#AP9976b). AP9976b was diluted at 1:100 dilution. An Alexa Fluor® 488-conjugated goat anti-mouse IgM at 1:400 dilution was used as the secondary antibody (green). DAPI was used to stain the cell nuclear (blue). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.